

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



B6

(43) International Publication Date
20 September 2001 (20.09.2001)

PCT

(10) International Publication Number
WO 01/68854 A2

(51) International Patent Classification⁷: C12N 15/12,
C07K 14/50, C12N 1/21, 5/10, C07K 16/22, A61K
39/395, G01N 33/68, A61K 38/18, 31/70, C12N 15/62,
A01K 67/027, C12N 5/22

(US). BASS, Michael, B. [US/US]; 1743 North Marian
Avenue, Thousand Oaks, CA 91360-2163 (US).

(21) International Application Number: PCT/US01/08013

(74) Agent: NOONAN, Kevin; McDonnell Boehnen Hulbert
& Berghoff, 300 South Wacker Drive, Suite 3200, Chicago,
IL 60606 (US).

(22) International Filing Date: 13 March 2001 (13.03.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/188,786 13 March 2000 (13.03.2000) US

(63) Related by continuation (CON) or continuation-in-part
(CIP) to earlier application:
US 60/188,786 (CON)
Filed on 13 March 2000 (13.03.2000)

(81) Designated States (*national*): AE, AL, AM, AT, AU, AZ,
BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE,
ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP,
KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD,
MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD,
SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ,
VN, YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SI, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,
CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant (*for all designated States except US*): AMGEN,
INC. [US/US]; One Amgen Center Drive, Thousand Oaks,
CA 91320-1799 (US).

Published:
— without international search report and to be republished
upon receipt of that report

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): JING, Shuqian
[CN/US]; 3254 Bordero Lane, Thousand Oaks, CA 91362

*For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.*

WO 01/68854 A2

(54) Title: FIBROBLAST GROWTH FACTOR-LIKE MOLECULES AND USES THEREOF

(57) Abstract: The present invention provides Fibroblast Growth Factor-Like (FGF-L) polypeptides and nucleic acid molecules encoding the same. The invention also provides selective binding agents, vectors, host cells, and methods for producing FGF-L polypeptides. The invention further provides pharmaceutical compositions and methods for the diagnosis, treatment, amelioration, and/or prevention of diseases, disorders, and conditions associated with FGF-L polypeptides.

BEST AVAILABLE COPY

FIBROBLAST GROWTH FACTOR-LIKE MOLECULES AND USES THEREOF

Field of the Invention

5 The present invention relates to Fibroblast Growth Factor-Like (FGF-L) polypeptides and nucleic acid molecules encoding the same. The invention also relates to selective binding agents, vectors, host cells, and methods for producing FGF-L polypeptides. The invention further relates to pharmaceutical compositions and methods for the diagnosis, treatment,
10 amelioration, and/or prevention of diseases, disorders, and conditions associated with FGF-L polypeptides.

Background of the Invention

Technical advances in the identification, cloning, expression, and
15 manipulation of nucleic acid molecules and the deciphering of the human genome have greatly accelerated the discovery of novel therapeutics. Rapid nucleic acid sequencing techniques can now generate sequence information at unprecedented rates and, coupled with computational analyses, allow the assembly of overlapping sequences into partial and entire genomes and the
20 identification of polypeptide-encoding regions. A comparison of a predicted amino acid sequence against a database compilation of known amino acid sequences allows one to determine the extent of homology to previously identified sequences and/or structural landmarks. The cloning and expression of a polypeptide-encoding region of a nucleic acid molecule provides a
25 polypeptide product for structural and functional analyses. The manipulation of nucleic acid molecules and encoded polypeptides may confer advantageous properties on a product for use as a therapeutic.

In spite of the significant technical advances in genome research over the past decade, the potential for the development of novel therapeutics based
30 on the human genome is still largely unrealized. Many genes encoding potentially beneficial polypeptide therapeutics or those encoding polypeptides, which may act as "targets" for therapeutic molecules, have still not been

identified. Accordingly, it is an object of the invention to identify novel polypeptides, and nucleic acid molecules encoding the same, which have diagnostic or therapeutic benefit.

5 Summary of the Invention

The present invention relates to novel FGF-L nucleic acid molecules and encoded polypeptides.

The invention provides for an isolated nucleic acid molecule comprising a nucleotide sequence selected from the group consisting of:

- 10 (a) the nucleotide sequence as set forth in SEQ ID NO: 1;
- (b) the nucleotide sequence of the DNA insert in ATCC Deposit No. PTA-1882;
- (c) a nucleotide sequence encoding the polypeptide as set forth in SEQ ID NO: 2;
- 15 (d) a nucleotide sequence which hybridizes under moderately or highly stringent conditions to the complement of any of (a) - (c); and
- (e) a nucleotide sequence complementary to any of (a) - (c).

The invention also provides for an isolated nucleic acid molecule
20 comprising a nucleotide sequence selected from the group consisting of:

- (a) a nucleotide sequence encoding a polypeptide which is at least about 70 percent identical to the polypeptide as set forth in SEQ ID NO: 2, wherein the encoded polypeptide has an activity of the polypeptide set forth in SEQ ID NO: 2;
- 25 (b) a nucleotide sequence encoding an allelic variant or splice variant of the nucleotide sequence as set forth in SEQ ID NO: 1, the nucleotide sequence of the DNA insert in ATCC Deposit No. PTA-1882, or (a);
- (c) a region of the nucleotide sequence of SEQ ID NO: 1, the DNA
30 insert in ATCC Deposit No. PTA-1882, (a), or (b) encoding a polypeptide fragment of at least about 25 amino acid residues, wherein the polypeptide

fragment has an activity of the polypeptide set forth in SEQ ID NO: 2, or is antigenic;

(d) a region of the nucleotide sequence of SEQ ID NO: 1, the DNA insert in ATCC Deposit No. PTA-1882, or any of (a) - (c) comprising a
5 fragment of at least about 16 nucleotides;

(e) a nucleotide sequence which hybridizes under moderately or highly stringent conditions to the complement of any of (a) - (d); and

(f) a nucleotide sequence complementary to any of (a) - (d).

10 The invention further provides for an isolated nucleic acid molecule comprising a nucleotide sequence selected from the group consisting of:

(a) a nucleotide sequence encoding a polypeptide as set forth in SEQ ID NO: 2 with at least one conservative amino acid substitution, wherein the encoded polypeptide has an activity of the polypeptide set forth in SEQ ID
15 NO: 2;

(b) a nucleotide sequence encoding a polypeptide as set forth in SEQ ID NO: 2 with at least one amino acid insertion, wherein the encoded polypeptide has an activity of the polypeptide set forth in SEQ ID NO: 2;

(c) a nucleotide sequence encoding a polypeptide as set forth in
20 SEQ ID NO: 2 with at least one amino acid deletion, wherein the encoded polypeptide has an activity of the polypeptide set forth in SEQ ID NO: 2;

(d) a nucleotide sequence encoding a polypeptide as set forth in SEQ ID NO: 2 which has a C- and/or N- terminal truncation, wherein the encoded polypeptide has an activity of the polypeptide set forth in SEQ ID
25 NO: 2;

(e) a nucleotide sequence encoding a polypeptide as set forth in SEQ ID NO: 2 with at least one modification selected from the group consisting of amino acid substitutions, amino acid insertions, amino acid deletions, C-terminal truncation, and N-terminal truncation, wherein the
30 encoded polypeptide has an activity of the polypeptide set forth in SEQ ID NO: 2;

(f) a nucleotide sequence of any of (a) - (e) comprising a fragment of at least about 16 nucleotides;

(g) a nucleotide sequence which hybridizes under moderately or highly stringent conditions to the complement of any of (a) - (f); and

5 (h) , a nucleotide sequence complementary to any of (a) - (e).

The present invention provides for an isolated polypeptide comprising an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence as set forth in SEQ ID NO: 2; and

10 (b) the amino acid sequence encoded by the DNA insert in ATCC Deposit No. PTA-1882.

The invention also provides for an isolated polypeptide comprising the amino acid sequence selected from the group consisting of:

15 (a) an amino acid sequence for an ortholog of SEQ ID NO: 2;

(b) an amino acid sequence which is at least about 70 percent identical to the amino acid sequence of SEQ ID NO: 2, wherein the polypeptide has an activity of the polypeptide set forth in SEQ ID NO: 2;

(c) a fragment of the amino acid sequence set forth in SEQ ID NO: 2 comprising at least about 25 amino acid residues, wherein the fragment has an activity of the polypeptide set forth in SEQ ID NO: 2, or is antigenic; and

20 (d) an amino acid sequence for an allelic variant or splice variant of the amino acid sequence as set forth in SEQ ID NO: 2, the amino acid sequence encoded by the DNA insert in ATCC Deposit No. PTA-1882, (a), or
25 (b).

The invention further provides for an isolated polypeptide comprising the amino acid sequence selected from the group consisting of:

(a) the amino acid sequence as set forth in SEQ ID NO: 2 with at
30 least one conservative amino acid substitution, wherein the polypeptide has an activity of the polypeptide set forth in SEQ ID NO: 2;

(b) the amino acid sequence as set forth in SEQ ID NO: 2 with at least one amino acid insertion, wherein the polypeptide has an activity of the polypeptide set forth in SEQ ID NO: 2;

(c) the amino acid sequence as set forth in SEQ ID NO: 2 with at least one amino acid deletion, wherein the polypeptide has an activity of the polypeptide set forth in SEQ ID NO: 2;

(d) the amino acid sequence as set forth in SEQ ID NO: 2 which has a C- and/or N-terminal truncation, wherein the polypeptide has an activity of the polypeptide set forth in SEQ ID NO: 2; and

(e) the amino acid sequence as set forth in SEQ ID NO: 2 with at least one modification selected from the group consisting of amino acid substitutions, amino acid insertions, amino acid deletions, C-terminal truncation, and N-terminal truncation, wherein the polypeptide has an activity of the polypeptide set forth in SEQ ID NO: 2.

15

Also provided are fusion polypeptides comprising FGF-L amino acid sequences.

The present invention also provides for an expression vector comprising the isolated nucleic acid molecules as set forth herein, recombinant host cells comprising the recombinant nucleic acid molecules as set forth herein, and a method of producing an FGF-L polypeptide comprising culturing the host cells and optionally isolating the polypeptide so produced.

A transgenic non-human animal comprising a nucleic acid molecule encoding an FGF-L polypeptide is also encompassed by the invention. The FGF-L nucleic acid molecules are introduced into the animal in a manner that allows expression and increased levels of an FGF-L polypeptide, which may include increased circulating levels. Alternatively, the FGF-L nucleic acid molecules are introduced into the animal in a manner that prevents expression of endogenous FGF-L polypeptide (*i.e.*, generates a transgenic animal possessing an FGF-L polypeptide gene knockout). The transgenic non-human animal is preferably a mammal, and more preferably a rodent, such as a rat or a mouse.

Also provided are derivatives of the FGF-L polypeptides of the present invention.

Additionally provided are selective binding agents such as antibodies and peptides capable of specifically binding the FGF-L polypeptides of the invention. Such antibodies and peptides may be agonistic or antagonistic.

Pharmaceutical compositions comprising the nucleotides, polypeptides, or selective binding agents of the invention and one or more pharmaceutically acceptable formulation agents are also encompassed by the invention. The pharmaceutical compositions are used to provide therapeutically effective amounts of the nucleotides or polypeptides of the present invention. The invention is also directed to methods of using the polypeptides, nucleic acid molecules, and selective binding agents.

The FGF-L polypeptides and nucleic acid molecules of the present invention may be used to treat, prevent, ameliorate, and/or detect diseases and disorders, including those recited herein.

The present invention also provides a method of assaying test molecules to identify a test molecule that binds to an FGF-L polypeptide. The method comprises contacting an FGF-L polypeptide with a test molecule to determine the extent of binding of the test molecule to the polypeptide. The method further comprises determining whether such test molecules are agonists or antagonists of an FGF-L polypeptide. The present invention further provides a method of testing the impact of molecules on the expression of FGF-L polypeptide or on the activity of FGF-L polypeptide.

Methods of regulating expression and modulating (*i.e.*, increasing or decreasing) levels of an FGF-L polypeptide are also encompassed by the invention. One method comprises administering to an animal a nucleic acid molecule encoding an FGF-L polypeptide. In another method, a nucleic acid molecule comprising elements that regulate or modulate the expression of an FGF-L polypeptide may be administered. Examples of these methods include gene therapy, cell therapy, and anti-sense therapy as further described herein.

In another aspect of the present invention, the FGF-L polypeptides may be used for identifying receptors thereof ("FGF-L polypeptide

receptors"). Various forms of "expression cloning" have been extensively used to clone receptors for protein ligands. *See, e.g.,* Simonsen and Lodish, 1994, *Trends Pharmacol. Sci.* 15:437-41 and Tartaglia *et al.*, 1995, *Cell* 83:1263-71. The isolation of an FGF-L polypeptide receptor is useful for
5 identifying or developing novel agonists and antagonists of the FGF-L polypeptide signaling pathway. Such agonists and antagonists include soluble FGF-L polypeptide receptors, anti-FGF-L polypeptide receptor-selective binding agents (such as antibodies and derivatives thereof), small molecules, and antisense oligonucleotides, any of which can be used for treating one or
10 more disease or disorder, including those disclosed herein.

Brief Description of the Figures

Figures 1A-1B illustrate the nucleotide sequence of the human FGF-L gene (SEQ ID NO: 1) and the deduced amino acid sequence of human FGF-L
15 polypeptide (SEQ ID NO: 2);

Figure 2 illustrates the amino acid sequence alignment of human FGF-L polypeptide (upper sequence; SEQ ID NO: 2) and human FGF-9 (lower
20 sequence; SEQ ID NO: 3);

Figure 3 illustrates the amino acid sequence alignment of human FGF-L polypeptide (upper sequence; SEQ ID NO: 2) and rat FGF-9 (lower sequence;
SEQ ID NO: 4);

25 Figure 4 illustrates the amino acid sequence alignment of human FGF-9 (hu FGF-9; SEQ ID NO: 3), human FGF-16 (hu FGF-16; SEQ ID NO: 5), human FGF-L polypeptide (hu FGF-L; SEQ ID NO: 2), murine FGF-9 (mu FGF-9; SEQ ID NO: 6), and rat FGF-16 (ra FGF-16; SEQ ID NO: 7).

30 Detailed Description of the Invention

The section headings used herein are for organizational purposes only and are not to be construed as limiting the subject matter described. All

references cited in this application are expressly incorporated by reference herein.

Definitions

5 The terms "FGF-L gene" or "FGF-L nucleic acid molecule" or "FGF-L polynucleotide" refer to a nucleic acid molecule comprising or consisting of a nucleotide sequence as set forth in SEQ ID NO: 1, a nucleotide sequence encoding the polypeptide as set forth in SEQ ID NO: 2, a nucleotide sequence of the DNA insert in ATCC Deposit No. PTA-1882, and nucleic acid
10 molecules as defined herein.

 The term "FGF-L polypeptide allelic variant" refers to one of several possible naturally occurring alternate forms of a gene occupying a given locus on a chromosome of an organism or a population of organisms.

 The term "FGF-L polypeptide splice variant" refers to a nucleic acid
15 molecule, usually RNA, which is generated by alternative processing of intron sequences in an RNA transcript of FGF-L polypeptide amino acid sequence as set forth in SEQ ID NO: 2.

 The term "isolated nucleic acid molecule" refers to a nucleic acid molecule of the invention that (1) has been separated from at least about 50
20 percent of proteins, lipids, carbohydrates, or other materials with which it is naturally found when total nucleic acid is isolated from the source cells, (2) is not linked to all or a portion of a polynucleotide to which the "isolated nucleic acid molecule" is linked in nature, (3) is operably linked to a polynucleotide which it is not linked to in nature, or (4) does not occur in nature as part of a
25 larger polynucleotide sequence. Preferably, the isolated nucleic acid molecule of the present invention is substantially free from any other contaminating nucleic acid molecule(s) or other contaminants that are found in its natural environment that would interfere with its use in polypeptide production or its therapeutic, diagnostic, prophylactic or research use.

30 The term "nucleic acid sequence" or "nucleic acid molecule" refers to a DNA or RNA sequence. The term encompasses molecules formed from any of the known base analogs of DNA and RNA such as, but not limited to 4-

acetylcytosine, 8-hydroxy-N6-methyladenosine, aziridinyl-cytosine, pseudoisocytosine, 5-(carboxyhydroxymethyl) uracil, 5-fluorouracil, 5-bromouracil, 5-carboxymethylaminomethyl-2-thiouracil, 5-carboxymethylaminomethyluracil, dihydrouracil, inosine, N6-iso-pentenyladenine, 1-methyladenine, 1-methylpseudouracil, 1-methylguanine, 1-methylinosine, 2,2-dimethyl-guanine, 2-methyladenine, 2-methylguanine, 3-methylcytosine, 5-methylcytosine, N6-methyladenine, 7-methylguanine, 5-methylaminomethyluracil, 5-methoxyamino-methyl-2-thiouracil, beta-D-mannosylqueosine, 5' -methoxycarbonyl-methyluracil, 5-methoxyuracil, 2-methylthio-N6-isopentenyladenine, uracil-5-oxyacetic acid methylester, uracil-5-oxyacetic acid, oxybutoxosine, pseudouracil, queosine, 2-thiocytosine, 5-methyl-2-thiouracil, 2-thiouracil, 4-thiouracil, 5-methyluracil, N-uracil-5-oxyacetic acid methylester, uracil-5-oxyacetic acid, pseudouracil, queosine, 2-thiocytosine, and 2,6-diaminopurine.

15 The term “vector” is used to refer to any molecule (*e.g.*, nucleic acid, plasmid, or virus) used to transfer coding information to a host cell.

The term “expression vector” refers to a vector that is suitable for transformation of a host cell and contains nucleic acid sequences that direct and/or control the expression of inserted heterologous nucleic acid sequences.

20 Expression includes, but is not limited to, processes such as transcription,
translation, and RNA splicing, if introns are present.

The term "operably linked" is used herein to refer to an arrangement of flanking sequences wherein the flanking sequences so described are configured or assembled so as to perform their usual function. Thus, a flanking sequence operably linked to a coding sequence may be capable of effecting the replication, transcription and/or translation of the coding sequence. For example, a coding sequence is operably linked to a promoter when the promoter is capable of directing transcription of that coding sequence. A flanking sequence need not be contiguous with the coding sequence, so long as it functions correctly. Thus, for example, intervening untranslated yet transcribed sequences can be present between a promoter

sequence and the coding sequence and the promoter sequence can still be considered "operably linked" to the coding sequence.

The term "host cell" is used to refer to a cell which has been transformed, or is capable of being transformed with a nucleic acid sequence
5 and then of expressing a selected gene of interest. The term includes the progeny of the parent cell, whether or not the progeny is identical in morphology or in genetic make-up to the original parent, so long as the selected gene is present.

The term "FGF-L polypeptide" refers to a polypeptide comprising the
10 amino acid sequence of SEQ ID NO: 2 and related polypeptides. Related polypeptides include FGF-L polypeptide fragments, FGF-L polypeptide orthologs, FGF-L polypeptide variants, and FGF-L polypeptide derivatives, which possess at least one activity of the polypeptide as set forth in SEQ ID NO: 2. FGF-L polypeptides may be mature polypeptides, as defined herein,
15 and may or may not have an amino-terminal methionine residue, depending on the method by which they are prepared.

The term "FGF-L polypeptide fragment" refers to a polypeptide that comprises a truncation at the amino-terminus (with or without a leader sequence) and/or a truncation at the carboxyl-terminus of the polypeptide as
20 set forth in SEQ ID NO: 2. The term "FGF-L polypeptide fragment" also refers to amino-terminal and/or carboxyl-terminal truncations of FGF-L polypeptide orthologs, FGF-L polypeptide derivatives, or FGF-L polypeptide variants, or to amino-terminal and/or carboxyl-terminal truncations of the polypeptides encoded by FGF-L polypeptide allelic variants or FGF-L
25 polypeptide splice variants. FGF-L polypeptide fragments may result from alternative RNA splicing or from *in vivo* protease activity. Membrane-bound forms of an FGF-L polypeptide are also contemplated by the present invention. In preferred embodiments, truncations and/or deletions comprise about 10 amino acids, or about 20 amino acids, or about 50 amino acids, or
30 about 75 amino acids, or about 100 amino acids, or more than about 100 amino acids. The polypeptide fragments so produced will comprise about 25 contiguous amino acids, or about 50 amino acids, or about 75 amino acids, or

about 100 amino acids, or about 150 amino acids, or more than about 150 amino acids. Such FGF-L polypeptide fragments may optionally comprise an amino-terminal methionine residue. It will be appreciated that such fragments can be used, for example, to generate antibodies to FGF-L polypeptides.

5 The term "FGF-L polypeptide ortholog" refers to a polypeptide from another species that corresponds to FGF-L polypeptide amino acid sequence as set forth in SEQ ID NO: 2. For example, mouse and human FGF-L polypeptides are considered orthologs of each other.

 The term "FGF-L polypeptide variants" refers to FGF-L polypeptides
10 comprising amino acid sequences having one or more amino acid sequence substitutions, deletions (such as internal deletions and/or FGF-L polypeptide fragments), and/or additions (such as internal additions and/or FGF-L fusion polypeptides) as compared to the FGF-L polypeptide amino acid sequence set forth in SEQ ID NO: 2 (with or without a leader sequence). Variants may be
15 naturally occurring (*e.g.*, FGF-L polypeptide allelic variants, FGF-L polypeptide orthologs, and FGF-L polypeptide splice variants) or artificially constructed. Such FGF-L polypeptide variants may be prepared from the corresponding nucleic acid molecules having a DNA sequence that varies accordingly from the DNA sequence as set forth in SEQ ID NO: 1. In
20 preferred embodiments, the variants have from 1 to 3, or from 1 to 5, or from 1 to 10, or from 1 to 15, or from 1 to 20, or from 1 to 25, or from 1 to 50, or from 1 to 75, or from 1 to 100, or more than 100 amino acid substitutions, insertions, additions and/or deletions, wherein the substitutions may be conservative, or non-conservative, or any combination thereof.

25 The term "FGF-L polypeptide derivatives" refers to the polypeptide as set forth in SEQ ID NO: 2, FGF-L polypeptide fragments, FGF-L polypeptide orthologs, or FGF-L polypeptide variants, as defined herein, that have been chemically modified. The term "FGF-L polypeptide derivatives" also refers to the polypeptides encoded by FGF-L polypeptide allelic variants or FGF-L
30 polypeptide splice variants, as defined herein, that have been chemically modified.

The term "mature FGF-L polypeptide" refers to an FGF-L polypeptide lacking a leader sequence. A mature FGF-L polypeptide may also include other modifications such as proteolytic processing of the amino-terminus (with or without a leader sequence) and/or the carboxyl-terminus, cleavage of a smaller polypeptide from a larger precursor, N-linked and/or O-linked glycosylation, and the like.

The term "FGF-L fusion polypeptide" refers to a fusion of one or more amino acids (such as a heterologous protein or peptide) at the amino- or carboxyl-terminus of the polypeptide as set forth in SEQ ID NO: 2, FGF-L polypeptide fragments, FGF-L polypeptide orthologs, FGF-L polypeptide variants, or FGF-L derivatives, as defined herein. The term "FGF-L fusion polypeptide" also refers to a fusion of one or more amino acids at the amino- or carboxyl-terminus of the polypeptide encoded by FGF-L polypeptide allelic variants or FGF-L polypeptide splice variants, as defined herein.

The term "biologically active FGF-L polypeptides" refers to FGF-L polypeptides having at least one activity characteristic of the polypeptide comprising the amino acid sequence of SEQ ID NO: 2. In addition, an FGF-L polypeptide may be active as an immunogen; that is, the FGF-L polypeptide contains at least one epitope to which antibodies may be raised.

The term "isolated polypeptide" refers to a polypeptide of the present invention that (1) has been separated from at least about 50 percent of polynucleotides, lipids, carbohydrates, or other materials with which it is naturally found when isolated from the source cell, (2) is not linked (by covalent or noncovalent interaction) to all or a portion of a polypeptide to which the "isolated polypeptide" is linked in nature, (3) is operably linked (by covalent or noncovalent interaction) to a polypeptide with which it is not linked in nature, or (4) does not occur in nature. Preferably, the isolated polypeptide is substantially free from any other contaminating polypeptides or other contaminants that are found in its natural environment that would interfere with its therapeutic, diagnostic, prophylactic or research use.

The term "identity," as known in the art, refers to a relationship between the sequences of two or more polypeptide molecules or two or more

nucleic acid molecules, as determined by comparing the sequences. In the art, "identity" also means the degree of sequence relatedness between nucleic acid molecules or polypeptides, as the case may be, as determined by the match between strings of two or more nucleotide or two or more amino acid sequences. "Identity" measures the percent of identical matches between the smaller of two or more sequences with gap alignments (if any) addressed by a particular mathematical model or computer program (*i.e.*, "algorithms").

The term "similarity" is a related concept, but in contrast to "identity," "similarity" refers to a measure of relatedness which includes both identical matches and conservative substitution matches. If two polypeptide sequences have, for example, 10/20 identical amino acids, and the remainder are all non-conservative substitutions, then the percent identity and similarity would both be 50%. If in the same example, there are five more positions where there are conservative substitutions, then the percent identity remains 50%, but the percent similarity would be 75% (15/20). Therefore, in cases where there are conservative substitutions, the percent similarity between two polypeptides will be higher than the percent identity between those two polypeptides.

The term "naturally occurring" or "native" when used in connection with biological materials such as nucleic acid molecules, polypeptides, host cells, and the like, refers to materials which are found in nature and are not manipulated by man. Similarly, "non-naturally occurring" or "non-native" as used herein refers to a material that is not found in nature or that has been structurally modified or synthesized by man.

The terms "effective amount" and "therapeutically effective amount" each refer to the amount of an FGF-L polypeptide or FGF-L nucleic acid molecule used to support an observable level of one or more biological activities of the FGF-L polypeptides as set forth herein.

The term "pharmaceutically acceptable carrier" or "physiologically acceptable carrier" as used herein refers to one or more formulation materials suitable for accomplishing or enhancing the delivery of the FGF-L polypeptide, FGF-L nucleic acid molecule, or FGF-L selective binding agent as a pharmaceutical composition.

The term "antigen" refers to a molecule or a portion of a molecule capable of being bound by a selective binding agent, such as an antibody, and additionally capable of being used in an animal to produce antibodies capable of binding to an epitope of that antigen. An antigen may have one or more
5 epitopes.

The term "selective binding agent" refers to a molecule or molecules having specificity for an FGF-L polypeptide. As used herein, the terms, "specific" and "specificity" refer to the ability of the selective binding agents to bind to human FGF-L polypeptides and not to bind to human non-FGF-L
10 polypeptides. It will be appreciated, however, that the selective binding agents may also bind orthologs of the polypeptide as set forth in SEQ ID NO: 2, that is, interspecies versions thereof, such as mouse and rat FGF-L polypeptides.

The term "transduction" is used to refer to the transfer of genes from one bacterium to another, usually by a phage. "Transduction" also refers to
15 the acquisition and transfer of eukaryotic cellular sequences by retroviruses.

The term "transfection" is used to refer to the uptake of foreign or exogenous DNA by a cell, and a cell has been "transfected" when the exogenous DNA has been introduced inside the cell membrane. A number of transfection techniques are well known in the art and are disclosed herein.
20 See, e.g., Graham *et al.*, 1973, *Virology* 52:456; Sambrook *et al.*, *Molecular Cloning, A Laboratory Manual* (Cold Spring Harbor Laboratories, 1989); Davis *et al.*, *Basic Methods in Molecular Biology* (Elsevier, 1986); and Chu *et al.*, 1981, *Gene* 13:197. Such techniques can be used to introduce one or more exogenous DNA moieties into suitable host cells.

The term "transformation" as used herein refers to a change in a cell's genetic characteristics, and a cell has been transformed when it has been modified to contain a new DNA. For example, a cell is transformed where it is genetically modified from its native state. Following transfection or transduction, the transforming DNA may recombine with that of the cell by
30 physically integrating into a chromosome of the cell, may be maintained transiently as an episomal element without being replicated, or may replicate

independently as a plasmid. A cell is considered to have been stably transformed when the DNA is replicated with the division of the cell.

Relatedness of Nucleic Acid Molecules and/or Polypeptides

5 It is understood that related nucleic acid molecules include allelic or splice variants of the nucleic acid molecule of SEQ ID NO: 1, and include sequences which are complementary to any of the above nucleotide sequences. Related nucleic acid molecules also include a nucleotide sequence encoding a polypeptide comprising or consisting essentially of a substitution,
10 modification, addition and/or deletion of one or more amino acid residues compared to the polypeptide in SEQ ID NO: 2. Such related FGF-L polypeptides may comprise, for example, an addition and/or a deletion of one or more N-linked or O-linked glycosylation sites or an addition and/or a deletion of one or more cysteine residues.

15 Related nucleic acid molecules also include fragments of FGF-L nucleic acid molecules which encode a polypeptide of at least about 25 contiguous amino acids, or about 50 amino acids, or about 75 amino acids, or about 100 amino acids, or about 150 amino acids, or more than about 150 amino acid residues of the FGF-L polypeptide of SEQ ID NO: 2.

20 In addition, related FGF-L nucleic acid molecules also include those molecules which comprise nucleotide sequences which hybridize under moderately or highly stringent conditions as defined herein with the fully complementary sequence of the FGF-L nucleic acid molecule of SEQ ID NO: 1, or of a molecule encoding a polypeptide, which polypeptide comprises the
25 amino acid sequence as shown in SEQ ID NO: 2, or of a nucleic acid fragment as defined herein, or of a nucleic acid fragment encoding a polypeptide as defined herein. Hybridization probes may be prepared using the FGF-L sequences provided herein to screen cDNA, genomic or synthetic DNA libraries for related sequences. Regions of the DNA and/or amino acid
30 sequence of FGF-L polypeptide that exhibit significant identity to known sequences are readily determined using sequence alignment algorithms as

described herein and those regions may be used to design probes for screening.

The term "highly stringent conditions" refers to those conditions that are designed to permit hybridization of DNA strands whose sequences are highly complementary, and to exclude hybridization of significantly mismatched DNAs. Hybridization stringency is principally determined by temperature, ionic strength, and the concentration of denaturing agents such as formamide. Examples of "highly stringent conditions" for hybridization and washing are 0.015 M sodium FGF-Loride, 0.0015 M sodium citrate at 65-68°C or 0.015 M sodium FGF-Loride, 0.0015 M sodium citrate, and 50% formamide at 42°C. See Sambrook, Fritsch & Maniatis, *Molecular Cloning: A Laboratory Manual* (2nd ed., Cold Spring Harbor Laboratory, 1989); Anderson *et al.*, *Nucleic Acid Hybridisation: A Practical Approach* Ch. 4 (IRL Press Limited).

More stringent conditions (such as higher temperature, lower ionic strength, higher formamide, or other denaturing agent) may also be used – however, the rate of hybridization will be affected. Other agents may be included in the hybridization and washing buffers for the purpose of reducing non-specific and/or background hybridization. Examples are 0.1% bovine serum albumin, 0.1% polyvinyl-pyrrolidone, 0.1% sodium pyrophosphate, 0.1% sodium dodecylsulfate, NaDodSO₄, (SDS), ficoll, Denhardt's solution, sonicated salmon sperm DNA (or another non-complementary DNA), and dextran sulfate, although other suitable agents can also be used. The concentration and types of these additives can be changed without substantially affecting the stringency of the hybridization conditions. Hybridization experiments are usually carried out at pH 6.8-7.4; however, at typical ionic strength conditions, the rate of hybridization is nearly independent of pH. See Anderson *et al.*, *Nucleic Acid Hybridisation: A Practical Approach* Ch. 4 (IRL Press Limited).

Factors affecting the stability of DNA duplex include base composition, length, and degree of base pair mismatch. Hybridization conditions can be adjusted by one skilled in the art in order to accommodate

these variables and allow DNAs of different sequence relatedness to form hybrids. The melting temperature of a perfectly matched DNA duplex can be estimated by the following equation:

$$T_m(^{\circ}\text{C}) = 81.5 + 16.6(\log[\text{Na}^+]) + 0.41(\% \text{G+C}) - 600/N - 0.72(\% \text{formamide})$$

where N is the length of the duplex formed, $[\text{Na}^+]$ is the molar concentration of the sodium ion in the hybridization or washing solution, %G+C is the percentage of (guanine+cytosine) bases in the hybrid. For imperfectly matched hybrids, the melting temperature is reduced by approximately 1°C for each 1% mismatch.

The term "moderately stringent conditions" refers to conditions under which a DNA duplex with a greater degree of base pair mismatching than could occur under "highly stringent conditions" is able to form. Examples of typical "moderately stringent conditions" are 0.015 M sodium FGF-Loride, 0.0015 M sodium citrate at 50-65°C or 0.015 M sodium FGF-Loride, 0.0015 M sodium citrate, and 20% formamide at 37-50°C. By way of example, "moderately stringent conditions" of 50°C in 0.015 M sodium ion will allow about a 21% mismatch.

It will be appreciated by those skilled in the art that there is no absolute distinction between "highly stringent conditions" and "moderately stringent conditions." For example, at 0.015 M sodium ion (no formamide), the melting temperature of perfectly matched long DNA is about 71°C. With a wash at 65°C (at the same ionic strength), this would allow for approximately a 6% mismatch. To capture more distantly related sequences, one skilled in the art can simply lower the temperature or raise the ionic strength.

A good estimate of the melting temperature in 1M NaCl* for oligonucleotide probes up to about 20nt is given by:

$$T_m = 2^{\circ}\text{C per A-T base pair} + 4^{\circ}\text{C per G-C base pair}$$

*The sodium ion concentration in 6X salt sodium citrate (SSC) is 1M. See Suggs *et al.*, *Developmental Biology Using Purified Genes* 683 (Brown and Fox, eds., 1981).

High stringency washing conditions for oligonucleotides are usually at a temperature of 0-5°C below the T_m of the oligonucleotide in 6X SSC, 0.1% SDS.

In another embodiment, related nucleic acid molecules comprise or
5 consist of a nucleotide sequence that is at least about 70 percent identical to the nucleotide sequence as shown in SEQ ID NO: 1, or comprise or consist essentially of a nucleotide sequence encoding a polypeptide that is at least about 70 percent identical to the polypeptide as set forth in SEQ ID NO: 2. In preferred embodiments, the nucleotide sequences are about 75 percent, or
10 about 80 percent, or about 85 percent, or about 90 percent, or about 95, 96, 97, 98, or 99 percent identical to the nucleotide sequence as shown in SEQ ID NO: 1, or the nucleotide sequences encode a polypeptide that is about 75 percent, or about 80 percent, or about 85 percent, or about 90 percent, or about 95, 96, 97, 98, or 99 percent identical to the polypeptide sequence as set forth
15 in SEQ ID NO: 2. Related nucleic acid molecules encode polypeptides possessing at least one activity of the polypeptide set forth in SEQ ID NO: 2.

Differences in the nucleic acid sequence may result in conservative and/or non-conservative modifications of the amino acid sequence relative to the amino acid sequence of SEQ ID NO: 2.

20 Conservative modifications to the amino acid sequence of SEQ ID NO: 2 (and the corresponding modifications to the encoding nucleotides) will produce a polypeptide having functional and chemical characteristics similar to those of FGF-L polypeptides. In contrast, substantial modifications in the functional and/or chemical characteristics of FGF-L polypeptides may be
25 accomplished by selecting substitutions in the amino acid sequence of SEQ ID NO: 2 that differ significantly in their effect on maintaining (a) the structure of the molecular backbone in the area of the substitution, for example, as a sheet or helical conformation, (b) the charge or hydrophobicity of the molecule at the target site, or (c) the bulk of the side chain.

30 For example, a "conservative amino acid substitution" may involve a substitution of a native amino acid residue with a nonnative residue such that there is little or no effect on the polarity or charge of the amino acid residue at

that position. Furthermore, any native residue in the polypeptide may also be substituted with alanine, as has been previously described for "alanine scanning mutagenesis."

Conservative amino acid substitutions also encompass non-naturally occurring amino acid residues that are typically incorporated by chemical peptide synthesis rather than by synthesis in biological systems. These include peptidomimetics, and other reversed or inverted forms of amino acid moieties.

Naturally occurring residues may be divided into classes based on common side chain properties:

- 1) hydrophobic: norleucine, Met, Ala, Val, Leu, Ile;
- 2) neutral hydrophilic: Cys, Ser, Thr;
- 3) acidic: Asp, Glu;
- 4) basic: Asn, Gln, His, Lys, Arg;
- 5) residues that influence chain orientation: Gly, Pro; and
- 6) aromatic: Trp, Tyr, Phe.

For example, non-conservative substitutions may involve the exchange of a member of one of these classes for a member from another class. Such substituted residues may be introduced into regions of the human FGF-L polypeptide that are homologous with non-human FGF-L polypeptides, or into the non-homologous regions of the molecule.

In making such changes, the hydropathic index of amino acids may be considered. Each amino acid has been assigned a hydropathic index on the basis of its hydrophobicity and charge characteristics. The hydropathic indices are: isoleucine (+4.5); valine (+4.2); leucine (+3.8); phenylalanine (+2.8); cysteine/cystine (+2.5); methionine (+1.9); alanine (+1.8); glycine (-0.4); threonine (-0.7); serine (-0.8); tryptophan (-0.9); tyrosine (-1.3); proline (-1.6); histidine (-3.2); glutamate (-3.5); glutamine (-3.5); aspartate (-3.5); asparagine (-3.5); lysine (-3.9); and arginine (-4.5).

The importance of the hydropathic amino acid index in conferring interactive biological function on a protein is generally understood in the art (Kyte *et al.*, 1982, *J. Mol. Biol.* 157:105-31). It is known that certain amino acids may be substituted for other amino acids having a similar hydropathic

index or score and still retain a similar biological activity. In making changes based upon the hydropathic index, the substitution of amino acids whose hydropathic indices are within ± 2 is preferred, those which are within ± 1 are particularly preferred, and those within ± 0.5 are even more particularly preferred.

It is also understood in the art that the substitution of like amino acids can be made effectively on the basis of hydrophilicity, particularly where the biologically functionally equivalent protein or peptide thereby created is intended for use in immunological embodiments, as in the present case. The greatest local average hydrophilicity of a protein, as governed by the hydrophilicity of its adjacent amino acids, correlates with its immunogenicity and antigenicity, *i.e.*, with a biological property of the protein.

The following hydrophilicity values have been assigned to these amino acid residues: arginine (+3.0); lysine (+3.0); aspartate (+3.0 \pm 1); glutamate (+3.0 \pm 1); serine (+0.3); asparagine (+0.2); glutamine (+0.2); glycine (0); threonine (-0.4); proline (-0.5 \pm 1); alanine (-0.5); histidine (-0.5); cysteine (-1.0); methionine (-1.3); valine (-1.5); leucine (-1.8); isoleucine (-1.8); tyrosine (-2.3); phenylalanine (-2.5); and tryptophan (-3.4). In making changes based upon similar hydrophilicity values, the substitution of amino acids whose hydrophilicity values are within ± 2 is preferred, those which are within ± 1 are particularly preferred, and those within ± 0.5 are even more particularly preferred. One may also identify epitopes from primary amino acid sequences on the basis of hydrophilicity. These regions are also referred to as "epitopic core regions."

Desired amino acid substitutions (whether conservative or non-conservative) can be determined by those skilled in the art at the time such substitutions are desired. For example, amino acid substitutions can be used to identify important residues of the FGF-L polypeptide, or to increase or decrease the affinity of the FGF-L polypeptides described herein. Exemplary amino acid substitutions are set forth in Table I.

Table I

<u>Amino Acid Substitutions</u>		
Original Residues	Exemplary Substitutions	Preferred Substitutions
Ala	Val, Leu, Ile	Val
Arg	Lys, Gln, Asn	Lys
Asn	Gln	Gln
Asp	Glu	Glu
Cys	Ser, Ala	Ser
Gln	Asn	Asn
Glu	Asp	Asp
Gly	Pro, Ala	Ala
His	Asn, Gln, Lys, Arg	Arg
Ile	Leu, Val, Met, Ala, Phe, Norleucine	Leu
Leu	Norleucine, Ile, Val, Met, Ala, Phe	Ile
Lys	Arg, 1,4 Diamino-butyric Acid, Gln, Asn	Arg
Met	Leu, Phe, Ile	Leu
Phe	Leu, Val, Ile, Ala, Tyr	Leu
Pro	Ala	Gly
Ser	Thr, Ala, Cys	Thr
Thr	Ser	Ser
Trp	Tyr, Phe	Tyr
Tyr	Trp, Phe, Thr, Ser	Phe
Val	Ile, Met, Leu, Phe, Ala, Norleucine	Leu

A skilled artisan will be able to determine suitable variants of the polypeptide as set forth in SEQ ID NO: 2 using well-known techniques. For

5 identifying suitable areas of the molecule that may be changed without

destroying biological activity, one skilled in the art may target areas not believed to be important for activity. For example, when similar polypeptides with similar activities from the same species or from other species are known, one skilled in the art may compare the amino acid sequence of an FGF-L polypeptide to such similar polypeptides. With such a comparison, one can identify residues and portions of the molecules that are conserved among similar polypeptides. It will be appreciated that changes in areas of the FGF-L molecule that are not conserved relative to such similar polypeptides would be less likely to adversely affect the biological activity and/or structure of an FGF-L polypeptide. One skilled in the art would also know that, even in relatively conserved regions, one may substitute chemically similar amino acids for the naturally occurring residues while retaining activity (conservative amino acid residue substitutions). Therefore, even areas that may be important for biological activity or for structure may be subject to conservative amino acid substitutions without destroying the biological activity or without adversely affecting the polypeptide structure.

Additionally, one skilled in the art can review structure-function studies identifying residues in similar polypeptides that are important for activity or structure. In view of such a comparison, one can predict the importance of amino acid residues in an FGF-L polypeptide that correspond to amino acid residues that are important for activity or structure in similar polypeptides. One skilled in the art may opt for chemically similar amino acid substitutions for such predicted important amino acid residues of FGF-L polypeptides.

One skilled in the art can also analyze the three-dimensional structure and amino acid sequence in relation to that structure in similar polypeptides. In view of such information, one skilled in the art may predict the alignment of amino acid residues of FGF-L polypeptide with respect to its three dimensional structure. One skilled in the art may choose not to make radical changes to amino acid residues predicted to be on the surface of the protein, since such residues may be involved in important interactions with other molecules. Moreover, one skilled in the art may generate test variants

containing a single amino acid substitution at each amino acid residue. The variants could be screened using activity assays known to those with skill in the art. Such variants could be used to gather information about suitable variants. For example, if one discovered that a change to a particular amino acid residue resulted in destroyed, undesirably reduced, or unsuitable activity, variants with such a change would be avoided. In other words, based on information gathered from such routine experiments, one skilled in the art can readily determine the amino acids where further substitutions should be avoided either alone or in combination with other mutations.

10 A number of scientific publications have been devoted to the prediction of secondary structure. See Moulton, 1996, *Curr. Opin. Biotechnol.* 7:422-27; Chou *et al.*, 1974, *Biochemistry* 13:222-45; Chou *et al.*, 1974, *Biochemistry* 113:211-22; Chou *et al.*, 1978, *Adv. Enzymol. Relat. Areas Mol. Biol.* 47:45-48; Chou *et al.*, 1978, *Ann. Rev. Biochem.* 47:251-276; and Chou
15 *et al.*, 1979, *Biophys. J.* 26:367-84. Moreover, computer programs are currently available to assist with predicting secondary structure. One method of predicting secondary structure is based upon homology modeling. For example, two polypeptides or proteins which have a sequence identity of greater than 30%, or similarity greater than 40%, often have similar structural
20 topologies. The recent growth of the protein structural database (PDB) has provided enhanced predictability of secondary structure, including the potential number of folds within the structure of a polypeptide or protein. See Holm *et al.*, 1999, *Nucleic Acids Res.* 27:244-47. It has been suggested that there are a limited number of folds in a given polypeptide or protein and that
25 once a critical number of structures have been resolved, structural prediction will become dramatically more accurate (Brenner *et al.*, 1997, *Curr. Opin. Struct. Biol.* 7:369-76).

Additional methods of predicting secondary structure include "threading" (Jones, 1997, *Curr. Opin. Struct. Biol.* 7:377-87; Sippl *et al.*,
30 1996, *Structure* 4:15-19), "profile analysis" (Bowie *et al.*, 1991, *Science*, 253:164-70; Gribskov *et al.*, 1990, *Methods Enzymol.* 183:146-59; Gribskov *et*

al., 1987, *Proc. Nat. Acad. Sci. U.S.A.* 84:4355-58), and "evolutionary linkage" (See Holm *et al.*, *supra*, and Brenner *et al.*, *supra*).

Preferred FGF-L polypeptide variants include glycosylation variants wherein the number and/or type of glycosylation sites have been altered compared to the amino acid sequence set forth in SEQ ID NO: 2. In one embodiment, FGF-L polypeptide variants comprise a greater or a lesser number of N-linked glycosylation sites than the amino acid sequence set forth in SEQ ID NO: 2. An N-linked glycosylation site is characterized by the sequence: Asn-X-Ser or Asn-X-Thr, wherein the amino acid residue designated as X may be any amino acid residue except proline. The substitution of amino acid residues to create this sequence provides a potential new site for the addition of an N-linked carbohydrate chain. Alternatively, substitutions that eliminate this sequence will remove an existing N-linked carbohydrate chain. Also provided is a rearrangement of N-linked carbohydrate chains wherein one or more N-linked glycosylation sites (typically those that are naturally occurring) are eliminated and one or more new N-linked sites are created. Additional preferred FGF-L variants include cysteine variants, wherein one or more cysteine residues are deleted or substituted with another amino acid (*e.g.*, serine) as compared to the amino acid sequence set forth in SEQ ID NO: 2. Cysteine variants are useful when FGF-L polypeptides must be refolded into a biologically active conformation such as after the isolation of insoluble inclusion bodies. Cysteine variants generally have fewer cysteine residues than the native protein, and typically have an even number to minimize interactions resulting from unpaired cysteines.

In other embodiments, related nucleic acid molecules comprise or consist of a nucleotide sequence encoding a polypeptide as set forth in SEQ ID NO: 2 with at least one amino acid insertion and wherein the polypeptide has an activity of the polypeptide set forth in SEQ ID NO: 2, or a nucleotide sequence encoding a polypeptide as set forth in SEQ ID NO: 2 with at least one amino acid deletion and wherein the polypeptide has an activity of the polypeptide set forth in SEQ ID NO: 2. Related nucleic acid molecules also

comprise or consist of a nucleotide sequence encoding a polypeptide as set forth in SEQ ID NO: 2 wherein the polypeptide has a carboxyl- and/or amino-terminal truncation and further wherein the polypeptide has an activity of the polypeptide set forth in SEQ ID NO: 2. Related nucleic acid molecules also
5 comprise or consist of a nucleotide sequence encoding a polypeptide as set forth in SEQ ID NO: 2 with at least one modification selected from the group consisting of amino acid substitutions, amino acid insertions, amino acid deletions, carboxyl-terminal truncations, and amino-terminal truncations and wherein the polypeptide has an activity of the polypeptide set forth in SEQ ID
10 NO: 2.

In addition, the polypeptide comprising the amino acid sequence of SEQ ID NO: 2, or other FGF-L polypeptide, may be fused to a homologous polypeptide to form a homodimer or to a heterologous polypeptide to form a heterodimer. Heterologous peptides and polypeptides include, but are not
15 limited to: an epitope to allow for the detection and/or isolation of an FGF-L fusion polypeptide; a transmembrane receptor protein or a portion thereof, such as an extracellular domain or a transmembrane and intracellular domain; a ligand or a portion thereof which binds to a transmembrane receptor protein; an enzyme or portion thereof which is catalytically active; a polypeptide or
20 peptide which promotes oligomerization, such as a leucine zipper domain; a polypeptide or peptide which increases stability, such as an immunoglobulin constant region; and a polypeptide which has a therapeutic activity different from the polypeptide comprising the amino acid sequence as set forth in SEQ ID NO: 2, or other FGF-L polypeptide.

25 Fusions can be made either at the amino-terminus or at the carboxyl-terminus of the polypeptide comprising the amino acid sequence set forth in SEQ ID NO: 2, or other FGF-L polypeptide. Fusions may be direct with no linker or adapter molecule or may be through a linker or adapter molecule. A linker or adapter molecule may be one or more amino acid residues, typically
30 from about 20 to about 50 amino acid residues. A linker or adapter molecule may also be designed with a cleavage site for a DNA restriction endonuclease or for a protease to allow for the separation of the fused moieties. It will be

appreciated that once constructed, the fusion polypeptides can be derivatized according to the methods described herein.

In a further embodiment of the invention, the polypeptide comprising the amino acid sequence of SEQ ID NO: 2, or other FGF-L polypeptide, is fused to one or more domains of an Fc region of human IgG. Antibodies comprise two functionally independent parts, a variable domain known as "Fab," that binds an antigen, and a constant domain known as "Fc," that is involved in effector functions such as complement activation and attack by phagocytic cells. An Fc has a long serum half-life, whereas an Fab is short-lived. Capon *et al.*, 1989, *Nature* 337:525-31. When constructed together with a therapeutic protein, an Fc domain can provide longer half-life or incorporate such functions as Fc receptor binding, protein A binding, complement fixation, and perhaps even placental transfer. *Id.* Table II summarizes the use of certain Fc fusions known in the art.

Table II

Fc Fusion with Therapeutic Proteins

Form of Fc	Fusion partner	Therapeutic implications	Reference
IgG1	N-terminus of CD30-L	Hodgkin's disease; anaplastic lymphoma; T-cell leukemia	U.S. Patent No. 5,480,981
Murine Fcγ2a	IL-10	anti-inflammatory; transplant rejection	Zheng <i>et al.</i> , 1995, <i>J. Immunol.</i> 154:5590-600
IgG1	TNF receptor	septic shock	Fisher <i>et al.</i> , 1996, <i>N. Engl. J. Med.</i> 334:1697-1702; Van Zee <i>et al.</i> , 1996, <i>J. Immunol.</i> 156:2221-30
IgG, IgA, IgM, or IgE (excluding the first domain)	TNF receptor	inflammation, autoimmune disorders	U.S. Patent No. 5,808,029
IgG1	CD4 receptor	AIDS	Capon <i>et al.</i> , 1989, <i>Nature</i> 337: 525-31
IgG1, IgG3	N-terminus of IL-2	anti-cancer, antiviral	Harvill <i>et al.</i> , 1995, <i>Immunotech.</i> 1:95-105
IgG1	C-terminus of OPG	osteoarthritis; bone density	WO 97/23614

IgG1	N-terminus of leptin	anti-obesity	PCT/US 97/23183, filed December 11, 1997
Human Ig C γ 1	CTLA-4	autoimmune disorders	Linsley, 1991, <i>J. Exp. Med.</i> , 174:561-69

In one example, a human IgG hinge, CH2, and CH3 region may be fused at either the amino-terminus or carboxyl-terminus of the FGF-L polypeptides using methods known to the skilled artisan. In another example,
5 a human IgG hinge, CH2, and CH3 region may be fused at either the amino-terminus or carboxyl-terminus of an FGF-L polypeptide fragment (*e.g.*, the predicted extracellular portion of FGF-L polypeptide).

The resulting FGF-L fusion polypeptide may be purified by use of a Protein A affinity column. Peptides and proteins fused to an Fc region have
10 been found to exhibit a substantially greater half-life *in vivo* than the unfused counterpart. Also, a fusion to an Fc region allows for dimerization/multimerization of the fusion polypeptide. The Fc region may be a naturally occurring Fc region, or may be altered to improve certain qualities, such as therapeutic qualities, circulation time, or reduced aggregation.

15 Identity and similarity of related nucleic acid molecules and polypeptides are readily calculated by known methods. Such methods include, but are not limited to those described in *Computational Molecular Biology* (A.M. Lesk, ed., Oxford University Press 1988); *Biocomputing: Informatics and Genome Projects* (D.W. Smith, ed., Academic Press 1993);
20 *Computer Analysis of Sequence Data* (Part 1, A.M. Griffin and H.G. Griffin, eds., Humana Press 1994); G. von Heinle, *Sequence Analysis in Molecular Biology* (Academic Press 1987); *Sequence Analysis Primer* (M. Gribskov and J. Devereux, eds., M. Stockton Press 1991); and Carillo *et al.*, 1988, *SIAM J. Applied Math.*, 48:1073.

25 Preferred methods to determine identity and/or similarity are designed to give the largest match between the sequences tested. Methods to determine identity and similarity are described in publicly available computer programs. Preferred computer program methods to determine identity and similarity between two sequences include, but are not limited to, the GCG program

package, including GAP (Devereux *et al.*, 1984, *Nucleic Acids Res.* 12:387; Genetics Computer Group, University of Wisconsin, Madison, WI), BLASTP, BLASTN, and FASTA (Altschul *et al.*, 1990, *J. Mol. Biol.* 215:403-10). The BLASTX program is publicly available from the National Center for
5 Biotechnology Information (NCBI) and other sources (Altschul *et al.*, *BLAST Manual* (NCB NLM NIH, Bethesda, MD); Altschul *et al.*, 1990, *supra*). The well-known Smith Waterman algorithm may also be used to determine identity.

Certain alignment schemes for aligning two amino acid sequences may
10 result in the matching of only a short region of the two sequences, and this small aligned region may have very high sequence identity even though there is no significant relationship between the two full-length sequences. Accordingly, in a preferred embodiment, the selected alignment method (GAP program) will result in an alignment that spans at least 50 contiguous amino
15 acids of the claimed polypeptide.

For example, using the computer algorithm GAP (Genetics Computer Group, University of Wisconsin, Madison, WI), two polypeptides for which the percent sequence identity is to be determined are aligned for optimal matching of their respective amino acids (the "matched span," as determined
20 by the algorithm). A gap opening penalty (which is calculated as 3X the average diagonal; the "average diagonal" is the average of the diagonal of the comparison matrix being used; the "diagonal" is the score or number assigned to each perfect amino acid match by the particular comparison matrix) and a gap extension penalty (which is usually 0.1X the gap opening penalty), as well
25 as a comparison matrix such as PAM 250 or BLOSUM 62 are used in conjunction with the algorithm. A standard comparison matrix is also used by the algorithm (see Dayhoff *et al.*, *5 Atlas of Protein Sequence and Structure* (Supp. 3 1978)(PAM250 comparison matrix); Henikoff *et al.*, 1992, *Proc. Natl. Acad. Sci USA* 89:10915-19 (BLOSUM 62 comparison matrix)).

30 Preferred parameters for polypeptide sequence comparison include the following:

Algorithm: Needleman and Wunsch, 1970, *J. Mol. Biol.* 48:443-53;

Comparison matrix: BLOSUM 62 (Henikoff *et al.*, *supra*);

Gap Penalty: 12

Gap Length Penalty: 4

5 Threshold of Similarity: 0

The GAP program is useful with the above parameters. The aforementioned parameters are the default parameters for polypeptide comparisons (along with no penalty for end gaps) using the GAP algorithm.

10 Preferred parameters for nucleic acid molecule sequence comparison include the following:

Algorithm: Needleman and Wunsch, *supra*;

Comparison matrix: matches = +10, mismatch = 0

15 Gap Penalty: 50

Gap Length Penalty: 3

The GAP program is also useful with the above parameters. The aforementioned parameters are the default parameters for nucleic acid
20 molecule comparisons.

Other exemplary algorithms, gap opening penalties, gap extension penalties, comparison matrices, and thresholds of similarity may be used, including those set forth in the Program Manual, Wisconsin Package, Version 9, September, 1997. The particular choices to be made will be apparent to
25 those of skill in the art and will depend on the specific comparison to be made, such as DNA-to-DNA, protein-to-protein, protein-to-DNA; and additionally, whether the comparison is between given pairs of sequences (in which case GAP or BestFit are generally preferred) or between one sequence and a large database of sequences (in which case FASTA or BLASTA are preferred).

30

Nucleic Acid Molecules

The nucleic acid molecules encoding a polypeptide comprising the amino acid sequence of an FGF-L polypeptide can readily be obtained in a variety of ways including, without limitation, chemical synthesis, cDNA or genomic library screening, expression library screening, and/or PCR
5 amplification of cDNA.

Recombinant DNA methods used herein are generally those set forth in Sambrook *et al.*, *Molecular Cloning: A Laboratory Manual* (Cold Spring Harbor Laboratory Press, 1989) and/or *Current Protocols in Molecular Biology* (Ausubel *et al.*, eds., Green Publishers Inc. and Wiley and Sons
10 1994). The invention provides for nucleic acid molecules as described herein and methods for obtaining such molecules.

Where a gene encoding the amino acid sequence of an FGF-L polypeptide has been identified from one species, all or a portion of that gene may be used as a probe to identify orthologs or related genes from the same
15 species. The probes or primers may be used to screen cDNA libraries from various tissue sources believed to express the FGF-L polypeptide. In addition, part or all of a nucleic acid molecule having the sequence as set forth in SEQ ID NO: 1 may be used to screen a genomic library to identify and isolate a gene encoding the amino acid sequence of an FGF-L polypeptide. Typically,
20 conditions of moderate or high stringency will be employed for screening to minimize the number of false positives obtained from the screening.

Nucleic acid molecules encoding the amino acid sequence of FGF-L polypeptides may also be identified by expression cloning which employs the detection of positive clones based upon a property of the expressed protein.
25 Typically, nucleic acid libraries are screened by the binding an antibody or other binding partner (*e.g.*, receptor or ligand) to cloned proteins that are expressed and displayed on a host cell surface. The antibody or binding partner is modified with a detectable label to identify those cells expressing the desired clone.

30 Recombinant expression techniques conducted in accordance with the descriptions set forth below may be followed to produce these polynucleotides and to express the encoded polypeptides. For example, by inserting a nucleic

acid sequence that encodes the amino acid sequence of an FGF-L polypeptide into an appropriate vector, one skilled in the art can readily produce large quantities of the desired nucleotide sequence. The sequences can then be used to generate detection probes or amplification primers. Alternatively, a polynucleotide encoding the amino acid sequence of an FGF-L polypeptide can be inserted into an expression vector. By introducing the expression vector into an appropriate host, the encoded FGF-L polypeptide may be produced in large amounts.

Another method for obtaining a suitable nucleic acid sequence is the polymerase chain reaction (PCR). In this method, cDNA is prepared from poly(A)+RNA or total RNA using the enzyme reverse transcriptase. Two primers, typically complementary to two separate regions of cDNA encoding the amino acid sequence of an FGF-L polypeptide, are then added to the cDNA along with a polymerase such as *Taq* polymerase, and the polymerase amplifies the cDNA region between the two primers.

Another means of preparing a nucleic acid molecule encoding the amino acid sequence of an FGF-L polypeptide is chemical synthesis using methods well known to the skilled artisan such as those described by Engels *et al.*, 1989, *Angew. Chem. Intl. Ed.* 28:716-34. These methods include, *inter alia*, the phosphotriester, phosphoramidite, and H-phosphonate methods for nucleic acid synthesis. A preferred method for such chemical synthesis is polymer-supported synthesis using standard phosphoramidite chemistry. Typically, the DNA encoding the amino acid sequence of an FGF-L polypeptide will be several hundred nucleotides in length. Nucleic acids larger than about 100 nucleotides can be synthesized as several fragments using these methods. The fragments can then be ligated together to form the full-length nucleotide sequence of an FGF-L gene. Usually, the DNA fragment encoding the amino-terminus of the polypeptide will have an ATG, which encodes a methionine residue. This methionine may or may not be present on the mature form of the FGF-L polypeptide, depending on whether the polypeptide produced in the host cell is designed to be secreted from that cell. Other methods known to the skilled artisan may be used as well.

In certain embodiments, nucleic acid variants contain codons which have been altered for optimal expression of an FGF-L polypeptide in a given host cell. Particular codon alterations will depend upon the FGF-L polypeptide and host cell selected for expression. Such "codon optimization" can be carried out by a variety of methods, for example, by selecting codons which are preferred for use in highly expressed genes in a given host cell. Computer algorithms which incorporate codon frequency tables such as "Eco_high.Cod" for codon preference of highly expressed bacterial genes may be used and are provided by the University of Wisconsin Package Version 9.0 (Genetics Computer Group, Madison, WI). Other useful codon frequency tables include "Celegans_high.cod," "Celegans_low.cod," "Drosophila_high.cod," "Human_high.cod," "Maize_high.cod," and "Yeast_high.cod."

In some cases, it may be desirable to prepare nucleic acid molecules encoding FGF-L polypeptide variants. Nucleic acid molecules encoding variants may be produced using site directed mutagenesis, PCR amplification, or other appropriate methods, where the primer(s) have the desired point mutations (see Sambrook *et al.*, *supra*, and Ausubel *et al.*, *supra*, for descriptions of mutagenesis techniques). Chemical synthesis using methods described by Engels *et al.*, *supra*, may also be used to prepare such variants. Other methods known to the skilled artisan may be used as well.

Vectors and Host Cells

A nucleic acid molecule encoding the amino acid sequence of an FGF-L polypeptide is inserted into an appropriate expression vector using standard ligation techniques. The vector is typically selected to be functional in the particular host cell employed (*i.e.*, the vector is compatible with the host cell machinery such that amplification of the gene and/or expression of the gene can occur). A nucleic acid molecule encoding the amino acid sequence of an FGF-L polypeptide may be amplified/expressed in prokaryotic, yeast, insect (baculovirus systems) and/or eukaryotic host cells. Selection of the host cell will depend in part on whether an FGF-L polypeptide is to be post-

translationally modified (*e.g.*, glycosylated and/or phosphorylated). If so, yeast, insect, or mammalian host cells are preferable. For a review of expression vectors, *see Meth. Enz.*, vol. 185 (D.V. Goeddel, ed., Academic Press 1990).

5 Typically, expression vectors used in any of the host cells will contain sequences for plasmid maintenance and for cloning and expression of exogenous nucleotide sequences. Such sequences, collectively referred to as "flanking sequences" in certain embodiments will typically include one or more of the following nucleotide sequences: a promoter, one or more enhancer
10 sequences, an origin of replication, a transcriptional termination sequence, a complete intron sequence containing a donor and acceptor splice site, a sequence encoding a leader sequence for polypeptide secretion, a ribosome binding site, a polyadenylation sequence, a polylinker region for inserting the nucleic acid encoding the polypeptide to be expressed, and a selectable marker
15 element. Each of these sequences is discussed below.

 Optionally, the vector may contain a "tag"-encoding sequence, *i.e.*, an oligonucleotide molecule located at the 5' or 3' end of the FGF-L polypeptide coding sequence; the oligonucleotide sequence encodes polyHis (such as hexaHis), or another "tag" such as FLAG, HA (hemagglutinin influenza virus),
20 or *myc* for which commercially available antibodies exist. This tag is typically fused to the polypeptide upon expression of the polypeptide, and can serve as a means for affinity purification of the FGF-L polypeptide from the host cell. Affinity purification can be accomplished, for example, by column chromatography using antibodies against the tag as an affinity matrix.
25 Optionally, the tag can subsequently be removed from the purified FGF-L polypeptide by various means such as using certain peptidases for cleavage.

 Flanking sequences may be homologous (*i.e.*, from the same species and/or strain as the host cell), heterologous (*i.e.*, from a species other than the host cell species or strain), hybrid (*i.e.*, a combination of flanking sequences
30 from more than one source), or synthetic, or the flanking sequences may be native sequences which normally function to regulate FGF-L polypeptide expression. As such, the source of a flanking sequence may be any

prokaryotic or eukaryotic organism, any vertebrate or invertebrate organism, or any plant, provided that the flanking sequence is functional in, and can be activated by, the host cell machinery.

Flanking sequences useful in the vectors of this invention may be
5 obtained by any of several methods well known in the art. Typically, flanking sequences useful herein – other than the FGF-L gene flanking sequences – will have been previously identified by mapping and/or by restriction endonuclease digestion and can thus be isolated from the proper tissue source using the appropriate restriction endonucleases. In some cases, the full
10 nucleotide sequence of a flanking sequence may be known. Here, the flanking sequence may be synthesized using the methods described herein for nucleic acid synthesis or cloning.

Where all or only a portion of the flanking sequence is known, it may be obtained using PCR and/or by screening a genomic library with a suitable
15 oligonucleotide and/or flanking sequence fragment from the same or another species. Where the flanking sequence is not known, a fragment of DNA containing a flanking sequence may be isolated from a larger piece of DNA that may contain, for example, a coding sequence or even another gene or genes. Isolation may be accomplished by restriction endonuclease digestion
20 to produce the proper DNA fragment followed by isolation using agarose gel purification, Qiagen® column chromatography (Chatsworth, CA), or other methods known to the skilled artisan. The selection of suitable enzymes to accomplish this purpose will be readily apparent to one of ordinary skill in the art.

25 An origin of replication is typically a part of those prokaryotic expression vectors purchased commercially, and the origin aids in the amplification of the vector in a host cell. Amplification of the vector to a certain copy number can, in some cases, be important for the optimal expression of an FGF-L polypeptide. If the vector of choice does not contain
30 an origin of replication site, one may be chemically synthesized based on a known sequence, and ligated into the vector. For example, the origin of replication from the plasmid pBR322 (New England Biolabs, Beverly, MA) is

suitable for most gram-negative bacteria and various origins (*e.g.*, SV40, polyoma, adenovirus, vesicular stomatitis virus (VSV), or papillomaviruses such as HPV or BPV) are useful for cloning vectors in mammalian cells. Generally, the origin of replication component is not needed for mammalian
5 expression vectors (for example, the SV40 origin is often used only because it contains the early promoter).

A transcription termination sequence is typically located 3' of the end of a polypeptide coding region and serves to terminate transcription. Usually, a transcription termination sequence in prokaryotic cells is a G-C rich
10 fragment followed by a poly-T sequence. While the sequence is easily cloned from a library or even purchased commercially as part of a vector, it can also be readily synthesized using methods for nucleic acid synthesis such as those described herein.

A selectable marker gene element encodes a protein necessary for the
15 survival and growth of a host cell grown in a selective culture medium. Typical selection marker genes encode proteins that (a) confer resistance to antibiotics or other toxins, *e.g.*, ampicillin, tetracycline, or kanamycin for prokaryotic host cells; (b) complement auxotrophic deficiencies of the cell; or (c) supply critical nutrients not available from complex media. Preferred
20 selectable markers are the kanamycin resistance gene, the ampicillin resistance gene, and the tetracycline resistance gene. A neomycin resistance gene may also be used for selection in prokaryotic and eukaryotic host cells.

Other selection genes may be used to amplify the gene that will be expressed. Amplification is the process wherein genes that are in greater
25 demand for the production of a protein critical for growth are reiterated in tandem within the chromosomes of successive generations of recombinant cells. Examples of suitable selectable markers for mammalian cells include dihydrofolate reductase (DHFR) and thymidine kinase. The mammalian cell transformants are placed under selection pressure wherein only the
30 transformants are uniquely adapted to survive by virtue of the selection gene present in the vector. Selection pressure is imposed by culturing the transformed cells under conditions in which the concentration of selection

agent in the medium is successively changed, thereby leading to the amplification of both the selection gene and the DNA that encodes an FGF-L polypeptide. As a result, increased quantities of FGF-L polypeptide are synthesized from the amplified DNA.

5 A ribosome binding site is usually necessary for translation initiation of mRNA and is characterized by a Shine-Dalgarno sequence (prokaryotes) or a Kozak sequence (eukaryotes). The element is typically located 3' to the promoter and 5' to the coding sequence of an FGF-L polypeptide to be expressed. The Shine-Dalgarno sequence is varied but is typically a
10 polypurine (*i.e.*, having a high A-G content). Many Shine-Dalgarno sequences have been identified, each of which can be readily synthesized using methods set forth herein and used in a prokaryotic vector.

 A leader, or signal, sequence may be used to direct an FGF-L polypeptide out of the host cell. Typically, a nucleotide sequence encoding
15 the signal sequence is positioned in the coding region of an FGF-L nucleic acid molecule, or directly at the 5' end of an FGF-L polypeptide coding region. Many signal sequences have been identified, and any of those that are functional in the selected host cell may be used in conjunction with an FGF-L nucleic acid molecule. Therefore, a signal sequence may be homologous
20 (naturally occurring) or heterologous to the FGF-L nucleic acid molecule. Additionally, a signal sequence may be chemically synthesized using methods described herein. In most cases, the secretion of an FGF-L polypeptide from the host cell via the presence of a signal peptide will result in the removal of the signal peptide from the secreted FGF-L polypeptide. The signal sequence
25 may be a component of the vector, or it may be a part of an FGF-L nucleic acid molecule that is inserted into the vector.

 Included within the scope of this invention is the use of either a nucleotide sequence encoding a native FGF-L polypeptide signal sequence joined to an FGF-L polypeptide coding region or a nucleotide sequence
30 encoding a heterologous signal sequence joined to an FGF-L polypeptide coding region. The heterologous signal sequence selected should be one that is recognized and processed, *i.e.*, cleaved by a signal peptidase, by the host

cell. For prokaryotic host cells that do not recognize and process the native FGF-L polypeptide signal sequence, the signal sequence is substituted by a prokaryotic signal sequence selected, for example, from the group of the alkaline phosphatase, penicillinase, or heat-stable enterotoxin II leaders. For yeast secretion, the native FGF-L polypeptide signal sequence may be substituted by the yeast invertase, alpha factor, or acid phosphatase leaders. In mammalian cell expression the native signal sequence is satisfactory, although other mammalian signal sequences may be suitable.

In some cases, such as where glycosylation is desired in a eukaryotic host cell expression system, one may manipulate the various presequences to improve glycosylation or yield. For example, one may alter the peptidase cleavage site of a particular signal peptide, or add pro-sequences, which also may affect glycosylation. The final protein product may have, in the -1 position (relative to the first amino acid of the mature protein) one or more additional amino acids incident to expression, which may not have been totally removed. For example, the final protein product may have one or two amino acid residues found in the peptidase cleavage site, attached to the amino-terminus. Alternatively, use of some enzyme cleavage sites may result in a slightly truncated form of the desired FGF-L polypeptide, if the enzyme cuts at such area within the mature polypeptide.

In many cases, transcription of a nucleic acid molecule is increased by the presence of one or more introns in the vector; this is particularly true where a polypeptide is produced in eukaryotic host cells, especially mammalian host cells. The introns used may be naturally occurring within the FGF-L gene especially where the gene used is a full-length genomic sequence or a fragment thereof. Where the intron is not naturally occurring within the gene (as for most cDNAs), the intron may be obtained from another source. The position of the intron with respect to flanking sequences and the FGF-L gene is generally important, as the intron must be transcribed to be effective. Thus, when an FGF-L cDNA molecule is being transcribed, the preferred position for the intron is 3' to the transcription start site and 5' to the poly-A transcription termination sequence. Preferably, the intron or introns will be

located on one side or the other (*i.e.*, 5' or 3') of the cDNA such that it does not interrupt the coding sequence. Any intron from any source, including viral, prokaryotic and eukaryotic (plant or animal) organisms, may be used to practice this invention, provided that it is compatible with the host cell into which it is inserted. Also included herein are synthetic introns. Optionally, more than one intron may be used in the vector.

The expression and cloning vectors of the present invention will typically contain a promoter that is recognized by the host organism and operably linked to the molecule encoding the FGF-L polypeptide. Promoters are untranscribed sequences located upstream (*i.e.*, 5') to the start codon of a structural gene (generally within about 100 to 1000 bp) that control the transcription of the structural gene. Promoters are conventionally grouped into one of two classes: inducible promoters and constitutive promoters. Inducible promoters initiate increased levels of transcription from DNA under their control in response to some change in culture conditions, such as the presence or absence of a nutrient or a change in temperature. Constitutive promoters, on the other hand, initiate continual gene product production; that is, there is little or no control over gene expression. A large number of promoters, recognized by a variety of potential host cells, are well known. A suitable promoter is operably linked to the DNA encoding FGF-L polypeptide by removing the promoter from the source DNA by restriction enzyme digestion and inserting the desired promoter sequence into the vector. The native FGF-L promoter sequence may be used to direct amplification and/or expression of an FGF-L nucleic acid molecule. A heterologous promoter is preferred, however, if it permits greater transcription and higher yields of the expressed protein as compared to the native promoter, and if it is compatible with the host cell system that has been selected for use.

Promoters suitable for use with prokaryotic hosts include the beta-lactamase and lactose promoter systems; alkaline phosphatase; a tryptophan (*trp*) promoter system; and hybrid promoters such as the *tac* promoter. Other known bacterial promoters are also suitable. Their sequences have been published, thereby enabling one skilled in the art to ligate them to the desired

DNA sequence, using linkers or adapters as needed to supply any useful restriction sites.

Suitable promoters for use with yeast hosts are also well known in the art. Yeast enhancers are advantageously used with yeast promoters. Suitable
5 promoters for use with mammalian host cells are well known and include, but are not limited to, those obtained from the genomes of viruses such as polyoma virus, fowlpox virus, adenovirus (such as Adenovirus 2), bovine papilloma virus, avian sarcoma virus, cytomegalovirus, retroviruses, hepatitis-B virus and most preferably Simian Virus 40 (SV40). Other suitable
10 mammalian promoters include heterologous mammalian promoters, for example, heat-shock promoters and the actin promoter.

Additional promoters which may be of interest in controlling FGF-L gene expression include, but are not limited to: the SV40 early promoter region (Bernoist and Chambon, 1981, *Nature* 290:304-10); the CMV
15 promoter; the promoter contained in the 3' long terminal repeat of Rous sarcoma virus (Yamamoto, *et al.*, 1980, *Cell* 22:787-97); the herpes thymidine kinase promoter (Wagner *et al.*, 1981, *Proc. Natl. Acad. Sci. U.S.A.* 78:1444-45); the regulatory sequences of the metallothioneine gene (Brinster *et al.*, 1982, *Nature* 296:39-42); prokaryotic expression vectors such as the beta-
20 lactamase promoter (Villa-Kamaroff *et al.*, 1978, *Proc. Natl. Acad. Sci. U.S.A.*, 75:3727-31); or the tac promoter (DeBoer *et al.*, 1983, *Proc. Natl. Acad. Sci. U.S.A.*, 80:21-25). Also of interest are the following animal transcriptional control regions, which exhibit tissue specificity and have been utilized in transgenic animals: the elastase I gene control region which is active in
25 pancreatic acinar cells (Swift *et al.*, 1984, *Cell* 38:639-46; Ornitz *et al.*, 1986, *Cold Spring Harbor Symp. Quant. Biol.* 50:399-409 (1986); MacDonald, 1987, *Hepatology* 7:425-515); the insulin gene control region which is active in pancreatic beta cells (Hanahan, 1985, *Nature* 315:115-22); the immunoglobulin gene control region which is active in lymphoid cells
30 (Grosschedl *et al.*, 1984, *Cell* 38:647-58; Adames *et al.*, 1985, *Nature* 318:533-38; Alexander *et al.*, 1987, *Mol. Cell. Biol.*, 7:1436-44); the mouse mammary tumor virus control region which is active in testicular, breast,

lymphoid and mast cells (Leder *et al.*, 1986, *Cell* 45:485-95); the albumin gene control region which is active in liver (Pinkert *et al.*, 1987, *Genes and Devel.* 1:268-76); the alpha-feto-protein gene control region which is active in liver (Krumlauf *et al.*, 1985, *Mol. Cell. Biol.*, 5:1639-48; Hammer *et al.*, 1987, *Science* 235:53-58); the alpha 1-antitrypsin gene control region which is active in the liver (Kelsey *et al.*, 1987, *Genes and Devel.* 1:161-71); the beta-globin gene control region which is active in myeloid cells (Mogram *et al.*, 1985, *Nature* 315:338-40; Kollias *et al.*, 1986, *Cell* 46:89-94); the myelin basic protein gene control region which is active in oligodendrocyte cells in the brain (Readhead *et al.*, 1987, *Cell* 48:703-12); the myosin light chain-2 gene control region which is active in skeletal muscle (Sani, 1985, *Nature* 314:283-86); and the gonadotropic releasing hormone gene control region which is active in the hypothalamus (Mason *et al.*, 1986, *Science* 234:1372-78).

An enhancer sequence may be inserted into the vector to increase the transcription of a DNA encoding an FGF-L polypeptide of the present invention by higher eukaryotes. Enhancers are cis-acting elements of DNA, usually about 10-300 bp in length, that act on the promoter to increase transcription. Enhancers are relatively orientation and position independent. They have been found 5' and 3' to the transcription unit. Several enhancer sequences available from mammalian genes are known (*e.g.*, globin, elastase, albumin, alpha-feto-protein and insulin). Typically, however, an enhancer from a virus will be used. The SV40 enhancer, the cytomegalovirus early promoter enhancer, the polyoma enhancer, and adenovirus enhancers are exemplary enhancing elements for the activation of eukaryotic promoters. While an enhancer may be spliced into the vector at a position 5' or 3' to an FGF-L nucleic acid molecule, it is typically located at a site 5' from the promoter.

Expression vectors of the invention may be constructed from a starting vector such as a commercially available vector. Such vectors may or may not contain all of the desired flanking sequences. Where one or more of the flanking sequences described herein are not already present in the vector, they may be individually obtained and ligated into the vector. Methods used for

obtaining each of the flanking sequences are well known to one skilled in the art.

Preferred vectors for practicing this invention are those which are compatible with bacterial, insect, and mammalian host cells. Such vectors include, *inter alia*, pCRII, pCR3, and pcDNA3.1 (Invitrogen, San Diego, CA), pBSII (Stratagene, La Jolla, CA), pET15 (Novagen, Madison, WI), pGEX (Pharmacia Biotech, Piscataway, NJ), pEGFP-N2 (Clontech, Palo Alto, CA), pETL (BlueBacII, Invitrogen), pDSR-alpha (PCT Pub. No. WO 90/14363) and pFastBacDual (Gibco-BRL, Grand Island, NY).

Additional suitable vectors include, but are not limited to, cosmids, plasmids, or modified viruses, but it will be appreciated that the vector system must be compatible with the selected host cell. Such vectors include, but are not limited to plasmids such as Bluescript[®] plasmid derivatives (a high copy number ColE1-based phagemid, Stratagene Cloning Systems, La Jolla CA), PCR cloning plasmids designed for cloning Taq-amplified PCR products (*e.g.*, TOPO[™] TA Cloning[®] Kit, PCR2.1[®] plasmid derivatives, Invitrogen, Carlsbad, CA), and mammalian, yeast or virus vectors such as a baculovirus expression system (pBacPAK plasmid derivatives, Clontech, Palo Alto, CA).

After the vector has been constructed and a nucleic acid molecule encoding an FGF-L polypeptide has been inserted into the proper site of the vector, the completed vector may be inserted into a suitable host cell for amplification and/or polypeptide expression. The transformation of an expression vector for an FGF-L polypeptide into a selected host cell may be accomplished by well known methods including methods such as transfection, infection, calcium FGF-Loride, electroporation, microinjection, lipofection, DEAE-dextran method, or other known techniques. The method selected will in part be a function of the type of host cell to be used. These methods and other suitable methods are well known to the skilled artisan, and are set forth, for example, in Sambrook *et al.*, *supra*.

Host cells may be prokaryotic host cells (such as *E. coli*) or eukaryotic host cells (such as a yeast, insect, or vertebrate cell). The host cell, when cultured under appropriate conditions, synthesizes an FGF-L polypeptide

which can subsequently be collected from the culture medium (if the host cell secretes it into the medium) or directly from the host cell producing it (if it is not secreted). The selection of an appropriate host cell will depend upon various factors, such as desired expression levels, polypeptide modifications
5 that are desirable or necessary for activity (such as glycosylation or phosphorylation) and ease of folding into a biologically active molecule.

A number of suitable host cells are known in the art and many are available from the American Type Culture Collection (ATCC), Manassas, VA. Examples include, but are not limited to, mammalian cells, such as Chinese
10 hamster ovary cells (CHO), CHO DHFR(-) cells (Urlaub *et al.*, 1980, *Proc. Natl. Acad. Sci. U.S.A.* 97:4216-20), human embryonic kidney (HEK) 293 or 293T cells, or 3T3 cells. The selection of suitable mammalian host cells and methods for transformation, culture, amplification, screening, product production, and purification are known in the art. Other suitable mammalian
15 cell lines, are the monkey COS-1 and COS-7 cell lines, and the CV-1 cell line. Further exemplary mammalian host cells include primate cell lines and rodent cell lines, including transformed cell lines. Normal diploid cells, cell strains derived from *in vitro* culture of primary tissue, as well as primary explants, are also suitable. Candidate cells may be genotypically deficient in the selection
20 gene, or may contain a dominantly acting selection gene. Other suitable mammalian cell lines include but are not limited to, mouse neuroblastoma N2A cells, HeLa, mouse L-929 cells, 3T3 lines derived from Swiss, Balb-c or NIH mice, BHK or HaK hamster cell lines. Each of these cell lines is known by and available to those skilled in the art of protein expression.

25 Similarly useful as host cells suitable for the present invention are bacterial cells. For example, the various strains of *E. coli* (e.g., HB101, DH5 α , DH10, and MC1061) are well-known as host cells in the field of biotechnology. Various strains of *B. subtilis*, *Pseudomonas spp.*, other *Bacillus spp.*, *Streptomyces spp.*, and the like may also be employed in this
30 method.

Many strains of yeast cells known to those skilled in the art are also available as host cells for the expression of the polypeptides of the present

invention. Preferred yeast cells include, for example, *Saccharomyces cerevisiae* and *Pichia pastoris*.

Additionally, where desired, insect cell systems may be utilized in the methods of the present invention. Such systems are described, for example, in
5 Kitts *et al.*, 1993, *Biotechniques*, 14:810-17; Lucklow, 1993, *Curr. Opin. Biotechnol.* 4:564-72; and Lucklow *et al.*, 1993, *J. Virol.*, 67:4566-79. Preferred insect cells are Sf-9 and Hi5 (Invitrogen).

One may also use transgenic animals to express glycosylated FGF-L polypeptides. For example, one may use a transgenic milk-producing animal
10 (a cow or goat, for example) and obtain the present glycosylated polypeptide in the animal milk. One may also use plants to produce FGF-L polypeptides, however, in general, the glycosylation occurring in plants is different from that produced in mammalian cells, and may result in a glycosylated product which is not suitable for human therapeutic use.

15

Polypeptide Production

Host cells comprising an FGF-L polypeptide expression vector may be cultured using standard media well known to the skilled artisan. The media will usually contain all nutrients necessary for the growth and survival of the
20 cells. Suitable media for culturing *E. coli* cells include, for example, Luria Broth (LB) and/or Terrific Broth (TB). Suitable media for culturing eukaryotic cells include Roswell Park Memorial Institute medium 1640 (RPMI 1640), Minimal Essential Medium (MEM) and/or Dulbecco's Modified Eagle Medium (DMEM), all of which may be supplemented with serum
25 and/or growth factors as necessary for the particular cell line being cultured. A suitable medium for insect cultures is Grace's medium supplemented with yeastolate, lactalbumin hydrolysate, and/or fetal calf serum as necessary.

Typically, an antibiotic or other compound useful for selective growth of transfected or transformed cells is added as a supplement to the media. The
30 compound to be used will be dictated by the selectable marker element present on the plasmid with which the host cell was transformed. For example, where the selectable marker element is kanamycin resistance, the compound added to

the culture medium will be kanamycin. Other compounds for selective growth include ampicillin, tetracycline, and neomycin.

The amount of an FGF-L polypeptide produced by a host cell can be evaluated using standard methods known in the art. Such methods include, without limitation, Western blot analysis, SDS-polyacrylamide gel electrophoresis, non-denaturing gel electrophoresis, High Performance Liquid Chromatography (HPLC) separation, immunoprecipitation, and/or activity assays such as DNA binding gel shift assays.

If an FGF-L polypeptide has been designed to be secreted from the host cells, the majority of polypeptide may be found in the cell culture medium. If however, the FGF-L polypeptide is not secreted from the host cells, it will be present in the cytoplasm and/or the nucleus (for eukaryotic host cells) or in the cytosol (for gram-negative bacteria host cells).

For an FGF-L polypeptide situated in the host cell cytoplasm and/or nucleus (for eukaryotic host cells) or in the cytosol (for bacterial host cells), the intracellular material (including inclusion bodies for gram-negative bacteria) can be extracted from the host cell using any standard technique known to the skilled artisan. For example, the host cells can be lysed to release the contents of the periplasm/cytoplasm by French press, homogenization, and/or sonication followed by centrifugation.

If an FGF-L polypeptide has formed inclusion bodies in the cytosol, the inclusion bodies can often bind to the inner and/or outer cellular membranes and thus will be found primarily in the pellet material after centrifugation. The pellet material can then be treated at pH extremes or with a chaotropic agent such as a detergent, guanidine, guanidine derivatives, urea, or urea derivatives in the presence of a reducing agent such as dithiothreitol at alkaline pH or tris carboxyethyl phosphine at acid pH to release, break apart, and solubilize the inclusion bodies. The solubilized FGF-L polypeptide can then be analyzed using gel electrophoresis, immunoprecipitation, or the like.

If it is desired to isolate the FGF-L polypeptide, isolation may be accomplished using standard methods such as those described herein and in Marston *et al.*, 1990, *Meth. Enz.*, 182:264-75.

In some cases, an FGF-L polypeptide may not be biologically active upon isolation. Various methods for "refolding" or converting the polypeptide to its tertiary structure and generating disulfide linkages can be used to restore biological activity. Such methods include exposing the solubilized polypeptide to a pH usually above 7 and in the presence of a particular concentration of a chaotrope. The selection of chaotrope is very similar to the choices used for inclusion body solubilization, but usually the chaotrope is used at a lower concentration and is not necessarily the same as chaotropes used for the solubilization. In most cases the refolding/oxidation solution will also contain a reducing agent or the reducing agent plus its oxidized form in a specific ratio to generate a particular redox potential allowing for disulfide shuffling to occur in the formation of the protein's cysteine bridges. Some of the commonly used redox couples include cysteine/cystamine, glutathione (GSH)/dithiobis GSH, cupric FGF-L-oxide, dithiothreitol (DTT)/dithiane DTT, and 2-2-mercaptoethanol(bME)/dithio-b(ME). In many instances, a cosolvent may be used or may be needed to increase the efficiency of the refolding, and the more common reagents used for this purpose include glycerol, polyethylene glycol of various molecular weights, arginine and the like.

If inclusion bodies are not formed to a significant degree upon expression of an FGF-L polypeptide, then the polypeptide will be found primarily in the supernatant after centrifugation of the cell homogenate. The polypeptide may be further isolated from the supernatant using methods such as those described herein.

The purification of an FGF-L polypeptide from solution can be accomplished using a variety of techniques. If the polypeptide has been synthesized such that it contains a tag such as Hexahistidine (FGF-L polypeptide/hexaHis) or other small peptide such as FLAG (Eastman Kodak Co., New Haven, CT) or *myc* (Invitrogen, Carlsbad, CA) at either its carboxyl- or amino-terminus, it may be purified in a one-step process by passing the solution through an affinity column where the column matrix has a high affinity for the tag.

For example, polyhistidine binds with great affinity and specificity to nickel. Thus, an affinity column of nickel (such as the Qiagen[®] nickel columns) can be used for purification of FGF-L polypeptide/polyHis. See, e.g., *Current Protocols in Molecular Biology* § 10.11.8 (Ausubel *et al.*, eds.,
5 Green Publishers Inc. and Wiley and Sons 1993).

Additionally, FGF-L polypeptides may be purified through the use of a monoclonal antibody that is capable of specifically recognizing and binding to an FGF-L polypeptide.

Other suitable procedures for purification include, without limitation,
10 affinity chromatography, immunoaffinity chromatography, ion exchange chromatography, molecular sieve chromatography, HPLC, electrophoresis (including native gel electrophoresis) followed by gel elution, and preparative isoelectric focusing ("Isoprime" machine/technique, Hoefer Scientific, San Francisco, CA). In some cases, two or more purification techniques may be
15 combined to achieve increased purity.

FGF-L polypeptides may also be prepared by chemical synthesis methods (such as solid phase peptide synthesis) using techniques known in the art such as those set forth by Merrifield *et al.*, 1963, *J. Am. Chem. Soc.* 85:2149; Houghten *et al.*, 1985, *Proc Natl Acad. Sci. USA* 82:5132; and
20 Stewart and Young, *Solid Phase Peptide Synthesis* (Pierce Chemical Co. 1984). Such polypeptides may be synthesized with or without a methionine on the amino-terminus. Chemically synthesized FGF-L polypeptides may be oxidized using methods set forth in these references to form disulfide bridges. Chemically synthesized FGF-L polypeptides are expected to have comparable
25 biological activity to the corresponding FGF-L polypeptides produced recombinantly or purified from natural sources, and thus may be used interchangeably with a recombinant or natural FGF-L polypeptide.

Another means of obtaining FGF-L polypeptide is via purification from biological samples such as source tissues and/or fluids in which the FGF-L
30 polypeptide is naturally found. Such purification can be conducted using methods for protein purification as described herein. The presence of the FGF-L polypeptide during purification may be monitored, for example, using

an antibody prepared against recombinantly produced FGF-L polypeptide or peptide fragments thereof.

A number of additional methods for producing nucleic acids and polypeptides are known in the art, and the methods can be used to produce polypeptides having specificity for FGF-L polypeptide. *See, e.g., Roberts et al., 1997, Proc. Natl. Acad. Sci. U.S.A. 94:12297-303*, which describes the production of fusion proteins between an mRNA and its encoded peptide. *See also, Roberts, 1999, Curr. Opin. Chem. Biol. 3:268-73*. Additionally, U.S. Patent No. 5,824,469 describes methods for obtaining oligonucleotides capable of carrying out a specific biological function. The procedure involves generating a heterogeneous pool of oligonucleotides, each having a 5' randomized sequence, a central preselected sequence, and a 3' randomized sequence. The resulting heterogeneous pool is introduced into a population of cells that do not exhibit the desired biological function. Subpopulations of the cells are then screened for those that exhibit a predetermined biological function. From that subpopulation, oligonucleotides capable of carrying out the desired biological function are isolated.

U.S. Patent Nos. 5,763,192; 5,814,476; 5,723,323; and 5,817,483 describe processes for producing peptides or polypeptides. This is done by producing stochastic genes or fragments thereof, and then introducing these genes into host cells which produce one or more proteins encoded by the stochastic genes. The host cells are then screened to identify those clones producing peptides or polypeptides having the desired activity.

Another method for producing peptides or polypeptides is described in PCT/US98/20094 (WO99/15650) filed by Athersys, Inc. Known as "Random Activation of Gene Expression for Gene Discovery" (RAGE-GD), the process involves the activation of endogenous gene expression or over-expression of a gene by *in situ* recombination methods. For example, expression of an endogenous gene is activated or increased by integrating a regulatory sequence into the target cell which is capable of activating expression of the gene by non-homologous or illegitimate recombination. The target DNA is first subjected to radiation, and a genetic promoter inserted. The promoter

eventually locates a break at the front of a gene, initiating transcription of the gene. This results in expression of the desired peptide or polypeptide.

It will be appreciated that these methods can also be used to create comprehensive FGF-L polypeptide expression libraries, which can subsequently be used for high throughput phenotypic screening in a variety of assays, such as biochemical assays, cellular assays, and whole organism assays (e.g., plant, mouse, etc.).

Synthesis

It will be appreciated by those skilled in the art that the nucleic acid and polypeptide molecules described herein may be produced by recombinant and other means.

Selective Binding Agents

The term "selective binding agent" refers to a molecule that has specificity for one or more FGF-L polypeptides. Suitable selective binding agents include, but are not limited to, antibodies and derivatives thereof, polypeptides, and small molecules. Suitable selective binding agents may be prepared using methods known in the art. An exemplary FGF-L polypeptide selective binding agent of the present invention is capable of binding a certain portion of the FGF-L polypeptide thereby inhibiting the binding of the polypeptide to an FGF-L polypeptide receptor.

Selective binding agents such as antibodies and antibody fragments that bind FGF-L polypeptides are within the scope of the present invention. The antibodies may be polyclonal including monospecific polyclonal; monoclonal (MAbs); recombinant; chimeric; humanized, such as CDR-grafted; human; single chain; and/or bispecific; as well as fragments; variants; or derivatives thereof. Antibody fragments include those portions of the antibody that bind to an epitope on the FGF-L polypeptide. Examples of such fragments include Fab and F(ab') fragments generated by enzymatic cleavage of full-length antibodies. Other binding fragments include those generated by recombinant DNA techniques, such as the expression of recombinant plasmids

containing nucleic acid sequences encoding antibody variable regions.

Polyclonal antibodies directed toward an FGF-L polypeptide generally are produced in animals (*e.g.*, rabbits or mice) by means of multiple subcutaneous or intraperitoneal injections of FGF-L polypeptide and an
5 adjuvant. It may be useful to conjugate an FGF-L polypeptide to a carrier protein that is immunogenic in the species to be immunized, such as keyhole limpet hemocyanin, serum, albumin, bovine thyroglobulin, or soybean trypsin inhibitor. Also, aggregating agents such as alum are used to enhance the immune response. After immunization, the animals are bled and the serum is
10 assayed for anti-FGF-L antibody titer.

Monoclonal antibodies directed toward FGF-L polypeptides are produced using any method that provides for the production of antibody molecules by continuous cell lines in culture. Examples of suitable methods for preparing monoclonal antibodies include the hybridoma methods of
15 Kohler *et al.*, 1975, *Nature* 256:495-97 and the human B-cell hybridoma method (Kozbor, 1984, *J. Immunol.* 133:3001; Brodeur *et al.*, *Monoclonal Antibody Production Techniques and Applications* 51-63 (Marcel Dekker, Inc., 1987). Also provided by the invention are hybridoma cell lines that produce monoclonal antibodies reactive with FGF-L polypeptides.

20 Monoclonal antibodies of the invention may be modified for use as therapeutics. One embodiment is a "chimeric" antibody in which a portion of the heavy (H) and/or light (L) chain is identical with or homologous to a corresponding sequence in antibodies derived from a particular species or belonging to a particular antibody class or subclass, while the remainder of the
25 chain(s) is/are identical with or homologous to a corresponding sequence in antibodies derived from another species or belonging to another antibody class or subclass. Also included are fragments of such antibodies, so long as they exhibit the desired biological activity. See U.S. Patent No. 4,816,567; Morrison *et al.*, 1985, *Proc. Natl. Acad. Sci.* 81:6851-55.

30 In another embodiment, a monoclonal antibody of the invention is a "humanized" antibody. Methods for humanizing non-human antibodies are well known in the art. See U.S. Patent Nos. 5,585,089 and 5,693,762.

Generally, a humanized antibody has one or more amino acid residues introduced into it from a source that is non-human. Humanization can be performed, for example, using methods described in the art (Jones *et al.*, 1986, *Nature* 321:522-25; Riechmann *et al.*, 1998, *Nature* 332:323-27; Verhoeyen *et al.*, 1988, *Science* 239:1534-36), by substituting at least a portion of a rodent complementarity-determining region (CDR) for the corresponding regions of a human antibody.

Also encompassed by the invention are human antibodies that bind FGF-L polypeptides. Using transgenic animals (*e.g.*, mice) that are capable of producing a repertoire of human antibodies in the absence of endogenous immunoglobulin production such antibodies are produced by immunization with an FGF-L polypeptide antigen (*i.e.*, having at least 6 contiguous amino acids), optionally conjugated to a carrier. *See, e.g.*, Jakobovits *et al.*, 1993, *Proc. Natl. Acad. Sci.* 90:2551-55; Jakobovits *et al.*, 1993, *Nature* 362:255-58; Bruggermann *et al.*, 1993, *Year in Immuno.* 7:33. In one method, such transgenic animals are produced by incapacitating the endogenous loci encoding the heavy and light immunoglobulin chains therein, and inserting loci encoding human heavy and light chain proteins into the genome thereof. Partially modified animals, that is those having less than the full complement of modifications, are then cross-bred to obtain an animal having all of the desired immune system modifications. When administered an immunogen, these transgenic animals produce antibodies with human (rather than, *e.g.*, murine) amino acid sequences, including variable regions which are immunospecific for these antigens. *See* PCT App. Nos. PCT/US96/05928 and PCT/US93/06926. Additional methods are described in U.S. Patent No. 5,545,807, PCT App. Nos. PCT/US91/245 and PCT/GB89/01207, and in European Patent Nos. 546073B1 and 546073A1. Human antibodies can also be produced by the expression of recombinant DNA in host cells or by expression in hybridoma cells as described herein.

In an alternative embodiment, human antibodies can also be produced from phage-display libraries (Hoogenboom *et al.*, 1991, *J. Mol. Biol.* 227:381; Marks *et al.*, 1991, *J. Mol. Biol.* 222:581). These processes mimic

immune selection through the display of antibody repertoires on the surface of filamentous bacteriophage, and subsequent selection of phage by their binding to an antigen of choice. One such technique is described in PCT App. No. PCT/US98/17364, which describes the isolation of high affinity and functional
5 agonistic antibodies for MPL- and msk- receptors using such an approach.

Chimeric, CDR grafted, and humanized antibodies are typically produced by recombinant methods. Nucleic acids encoding the antibodies are introduced into host cells and expressed using materials and procedures described herein. In a preferred embodiment, the antibodies are produced in
10 mammalian host cells, such as CHO cells. Monoclonal (e.g., human) antibodies may be produced by the expression of recombinant DNA in host cells or by expression in hybridoma cells as described herein.

The anti-FGF-L antibodies of the invention may be employed in any known assay method, such as competitive binding assays, direct and indirect
15 sandwich assays, and immunoprecipitation assays (Sola, *Monoclonal Antibodies: A Manual of Techniques* 147-158 (CRC Press, Inc., 1987)) for the detection and quantitation of FGF-L polypeptides. The antibodies will bind FGF-L polypeptides with an affinity that is appropriate for the assay method being employed.

20 For diagnostic applications, in certain embodiments, anti-FGF-L antibodies may be labeled with a detectable moiety. The detectable moiety can be any one that is capable of producing, either directly or indirectly, a detectable signal. For example, the detectable moiety may be a radioisotope, such as ^3H , ^{14}C , ^{32}P , ^{35}S , ^{125}I , ^{99}Tc , ^{111}In , or ^{67}Ga ; a fluorescent or
25 chemiluminescent compound, such as fluorescein isothiocyanate, rhodamine, or luciferin; or an enzyme, such as alkaline phosphatase, β -galactosidase, or horseradish peroxidase (Bayer, *et al.*, 1990, *Meth. Enz.* 184:138-63).

Competitive binding assays rely on the ability of a labeled standard (e.g., an FGF-L polypeptide, or an immunologically reactive portion thereof)
30 to compete with the test sample analyte (an FGF-L polypeptide) for binding with a limited amount of anti-FGF-L antibody. The amount of an FGF-L polypeptide in the test sample is inversely proportional to the amount of

standard that becomes bound to the antibodies. To facilitate determining the amount of standard that becomes bound, the antibodies typically are insolubilized before or after the competition, so that the standard and analyte that are bound to the antibodies may conveniently be separated from the
5 standard and analyte which remain unbound.

Sandwich assays typically involve the use of two antibodies, each capable of binding to a different immunogenic portion, or epitope, of the protein to be detected and/or quantitated. In a sandwich assay, the test sample analyte is typically bound by a first antibody which is immobilized on a solid
10 support, and thereafter a second antibody binds to the analyte, thus forming an insoluble three-part complex. See, e.g., U.S. Patent No. 4,376,110. The second antibody may itself be labeled with a detectable moiety (direct sandwich assays) or may be measured using an anti-immunoglobulin antibody that is labeled with a detectable moiety (indirect sandwich assays). For
15 example, one type of sandwich assay is an enzyme-linked immunosorbent assay (ELISA), in which case the detectable moiety is an enzyme.

The selective binding agents, including anti-FGF-L antibodies, are also useful for *in vivo* imaging. An antibody labeled with a detectable moiety may be administered to an animal, preferably into the bloodstream, and the
20 presence and location of the labeled antibody in the host assayed. The antibody may be labeled with any moiety that is detectable in an animal, whether by nuclear magnetic resonance, radiology, or other detection means known in the art.

Selective binding agents of the invention, including antibodies, may be
25 used as therapeutics. These therapeutic agents are generally agonists or antagonists, in that they either enhance or reduce, respectively, at least one of the biological activities of an FGF-L polypeptide. In one embodiment, antagonist antibodies of the invention are antibodies or binding fragments thereof which are capable of specifically binding to an FGF-L polypeptide and
30 which are capable of inhibiting or eliminating the functional activity of an FGF-L polypeptide *in vivo* or *in vitro*. In preferred embodiments, the selective binding agent, e.g., an antagonist antibody, will inhibit the functional activity

of an FGF-L polypeptide by at least about 50%, and preferably by at least about 80%. In another embodiment, the selective binding agent may be an anti-FGF-L polypeptide antibody that is capable of interacting with an FGF-L polypeptide binding partner (a ligand or receptor) thereby inhibiting or
5 eliminating FGF-L polypeptide activity *in vitro* or *in vivo*. Selective binding agents, including agonist and antagonist anti-FGF-L polypeptide antibodies, are identified by screening assays that are well known in the art.

The invention also relates to a kit comprising FGF-L selective binding agents (such as antibodies) and other reagents useful for detecting FGF-L
10 polypeptide levels in biological samples. Such reagents may include a detectable label, blocking serum, positive and negative control samples, and detection reagents.

Microarrays

15 It will be appreciated that DNA microarray technology can be utilized in accordance with the present invention. DNA microarrays are miniature, high-density arrays of nucleic acids positioned on a solid support, such as glass. Each cell or element within the array contains numerous copies of a single nucleic acid species that acts as a target for hybridization with a
20 complementary nucleic acid sequence (*e.g.*, mRNA). In expression profiling using DNA microarray technology, mRNA is first extracted from a cell or tissue sample and then converted enzymatically to fluorescently labeled cDNA. This material is hybridized to the microarray and unbound cDNA is removed by washing. The expression of discrete genes represented on the
25 array is then visualized by quantitating the amount of labeled cDNA that is specifically bound to each target nucleic acid molecule. In this way, the expression of thousands of genes can be quantitated in a high throughput, parallel manner from a single sample of biological material.

This high throughput expression profiling has a broad range of
30 applications with respect to the FGF-L molecules of the invention, including, but not limited to: the identification and validation of FGF-L disease-related genes as targets for therapeutics; molecular toxicology of related FGF-L

molecules and inhibitors thereof; stratification of populations and generation of surrogate markers for clinical trials; and enhancing related FGF-L polypeptide small molecule drug discovery by aiding in the identification of selective compounds in high throughput screens.

5

Chemical Derivatives

Chemically modified derivatives of FGF-L polypeptides may be prepared by one skilled in the art, given the disclosures described herein. FGF-L polypeptide derivatives are modified in a manner that is different –
10 either in the type or location of the molecules naturally attached to the polypeptide. Derivatives may include molecules formed by the deletion of one or more naturally-attached chemical groups. The polypeptide comprising the amino acid sequence of SEQ ID NO: 2, or other FGF-L polypeptide, may be modified by the covalent attachment of one or more polymers. For
15 example, the polymer selected is typically water-soluble so that the protein to which it is attached does not precipitate in an aqueous environment, such as a physiological environment. Included within the scope of suitable polymers is a mixture of polymers. Preferably, for therapeutic use of the end-product preparation, the polymer will be pharmaceutically acceptable.

20 The polymers each may be of any molecular weight and may be branched or unbranched. The polymers each typically have an average molecular weight of between about 2 kDa to about 100 kDa (the term “about” indicating that in preparations of a water-soluble polymer, some molecules will weigh more, some less, than the stated molecular weight). The average
25 molecular weight of each polymer is preferably between about 5 kDa and about 50 kDa, more preferably between about 12 kDa and about 40 kDa and most preferably between about 20 kDa and about 35 kDa.

Suitable water-soluble polymers or mixtures thereof include, but are not limited to, N-linked or O-linked carbohydrates, sugars, phosphates,
30 polyethylene glycol (PEG) (including the forms of PEG that have been used to derivatize proteins, including mono-(C₁-C₁₀), alkoxy-, or aryloxy-polyethylene glycol), monomethoxy-polyethylene glycol, dextran (such as low molecular

weight dextran of, for example, about 6 kD), cellulose, or other carbohydrate based polymers, poly-(N-vinyl pyrrolidone) polyethylene glycol, propylene glycol homopolymers, polypropylene oxide/ethylene oxide co-polymers, polyoxyethylated polyols (e.g., glycerol), and polyvinyl alcohol. Also
5 encompassed by the present invention are bifunctional crosslinking molecules which may be used to prepare covalently attached FGF-L polypeptide multimers.

In general, chemical derivatization may be performed under any suitable condition used to react a protein with an activated polymer molecule.
10 Methods for preparing chemical derivatives of polypeptides will generally comprise the steps of: (a) reacting the polypeptide with the activated polymer molecule (such as a reactive ester or aldehyde derivative of the polymer molecule) under conditions whereby the polypeptide comprising the amino acid sequence of SEQ ID NO: 2, or other FGF-L polypeptide, becomes
15 attached to one or more polymer molecules, and (b) obtaining the reaction products. The optimal reaction conditions will be determined based on known parameters and the desired result. For example, the larger the ratio of polymer molecules to protein, the greater the percentage of attached polymer molecule. In one embodiment, the FGF-L polypeptide derivative may have a single
20 polymer molecule moiety at the amino-terminus. *See, e.g.,* U.S. Patent No. 5,234,784.

The pegylation of a polypeptide may be specifically carried out using any of the pegylation reactions known in the art. Such reactions are described, for example, in the following references: Francis *et al.*, 1992, *Focus on*
25 *Growth Factors* 3:4-10; European Patent Nos. 0154316 and 0401384; and U.S. Patent No. 4,179,337. For example, pegylation may be carried out via an acylation reaction or an alkylation reaction with a reactive polyethylene glycol molecule (or an analogous reactive water-soluble polymer) as described herein. For the acylation reactions, a selected polymer should have a single
30 reactive ester group. For reductive alkylation, a selected polymer should have a single reactive aldehyde group. A reactive aldehyde is, for example,

polyethylene glycol propionaldehyde, which is water stable, or mono C₁-C₁₀ alkoxy or aryloxy derivatives thereof (*see* U.S. Patent No. 5,252,714).

In another embodiment, FGF-L polypeptides may be chemically coupled to biotin. The biotin/FGF-L polypeptide molecules are then allowed
5 to bind to avidin, resulting in tetravalent avidin/biotin/FGF-L polypeptide molecules. FGF-L polypeptides may also be covalently coupled to dinitrophenol (DNP) or trinitrophenol (TNP) and the resulting conjugates precipitated with anti-DNP or anti-TNP-IgM to form decameric conjugates with a valency of 10.

10 Generally, conditions that may be alleviated or modulated by the administration of the present FGF-L polypeptide derivatives include those described herein for FGF-L polypeptides. However, the FGF-L polypeptide derivatives disclosed herein may have additional activities, enhanced or reduced biological activity, or other characteristics, such as increased or
15 decreased half-life, as compared to the non-derivatized molecules.

Genetically Engineered Non-Human Animals

Additionally included within the scope of the present invention are non-human animals such as mice, rats, or other rodents; rabbits, goats, sheep,
20 or other farm animals, in which the genes encoding native FGF-L polypeptide have been disrupted (*i.e.*, "knocked out") such that the level of expression of FGF-L polypeptide is significantly decreased or completely abolished. Such animals may be prepared using techniques and methods such as those described in U.S. Patent No. 5,557,032.

25 The present invention further includes non-human animals such as mice, rats, or other rodents; rabbits, goats, sheep, or other farm animals, in which either the native form of an FGF-L gene for that animal or a heterologous FGF-L gene is over-expressed by the animal, thereby creating a "transgenic" animal. Such transgenic animals may be prepared using well
30 known methods such as those described in U.S. Patent No 5,489,743 and PCT Pub. No. WO 94/28122.

The present invention further includes non-human animals in which the promoter for one or more of the FGF-L polypeptides of the present invention is either activated or inactivated (e.g., by using homologous recombination methods) to alter the level of expression of one or more of the native FGF-L polypeptides.

These non-human animals may be used for drug candidate screening. In such screening, the impact of a drug candidate on the animal may be measured. For example, drug candidates may decrease or increase the expression of the FGF-L gene. In certain embodiments, the amount of FGF-L polypeptide that is produced may be measured after the exposure of the animal to the drug candidate. Additionally, in certain embodiments, one may detect the actual impact of the drug candidate on the animal. For example, over-expression of a particular gene may result in, or be associated with, a disease or pathological condition. In such cases, one may test a drug candidate's ability to decrease expression of the gene or its ability to prevent or inhibit a pathological condition. In other examples, the production of a particular metabolic product such as a fragment of a polypeptide, may result in, or be associated with, a disease or pathological condition. In such cases, one may test a drug candidate's ability to decrease the production of such a metabolic product or its ability to prevent or inhibit a pathological condition.

Assaying for Other Modulators of FGF-L Polypeptide Activity

In some situations, it may be desirable to identify molecules that are modulators, *i.e.*, agonists or antagonists, of the activity of FGF-L polypeptide. Natural or synthetic molecules that modulate FGF-L polypeptide may be identified using one or more screening assays, such as those described herein. Such molecules may be administered either in an *ex vivo* manner or in an *in vivo* manner by injection, or by oral delivery, implantation device, or the like.

"Test molecule" refers to a molecule that is under evaluation for the ability to modulate (*i.e.*, increase or decrease) the activity of an FGF-L polypeptide. Most commonly, a test molecule will interact directly with an FGF-L polypeptide. However, it is also contemplated that a test molecule

may also modulate FGF-L polypeptide activity indirectly, such as by affecting FGF-L gene expression, or by binding to an FGF-L polypeptide binding partner (*e.g.*, receptor or ligand). In one embodiment, a test molecule will bind to an FGF-L polypeptide with an affinity constant of at least about 10^{-6} M, preferably about 10^{-8} M, more preferably about 10^{-9} M, and even more preferably about 10^{-10} M.

Methods for identifying compounds that interact with FGF-L polypeptides are encompassed by the present invention. In certain embodiments, an FGF-L polypeptide is incubated with a test molecule under conditions that permit the interaction of the test molecule with an FGF-L polypeptide, and the extent of the interaction is measured. The test molecule can be screened in a substantially purified form or in a crude mixture.

In certain embodiments, an FGF-L polypeptide agonist or antagonist may be a protein, peptide, carbohydrate, lipid, or small molecular weight molecule that interacts with FGF-L polypeptide to regulate its activity. Molecules which regulate FGF-L polypeptide expression include nucleic acids which are complementary to nucleic acids encoding an FGF-L polypeptide, or are complementary to nucleic acids sequences which direct or control the expression of FGF-L polypeptide, and which act as anti-sense regulators of expression.

Once a test molecule has been identified as interacting with an FGF-L polypeptide, the molecule may be further evaluated for its ability to increase or decrease FGF-L polypeptide activity. The measurement of the interaction of a test molecule with FGF-L polypeptide may be carried out in several formats, including cell-based binding assays, membrane binding assays, solution-phase assays, and immunoassays. In general, a test molecule is incubated with an FGF-L polypeptide for a specified period of time, and FGF-L polypeptide activity is determined by one or more assays for measuring biological activity.

The interaction of test molecules with FGF-L polypeptides may also be assayed directly using polyclonal or monoclonal antibodies in an immunoassay. Alternatively, modified forms of FGF-L polypeptides

containing epitope tags as described herein may be used in solution and immunoassays.

In the event that FGF-L polypeptides display biological activity through an interaction with a binding partner (*e.g.*, a receptor or a ligand), a variety of *in vitro* assays may be used to measure the binding of an FGF-L polypeptide to the corresponding binding partner (such as a selective binding agent, receptor, or ligand). These assays may be used to screen test molecules for their ability to increase or decrease the rate and/or the extent of binding of an FGF-L polypeptide to its binding partner. In one assay, an FGF-L polypeptide is immobilized in the wells of a microtiter plate. Radiolabeled FGF-L polypeptide binding partner (for example, iodinated FGF-L polypeptide binding partner) and a test molecule can then be added either one at a time (in either order) or simultaneously to the wells. After incubation, the wells can be washed and counted for radioactivity, using a scintillation counter, to determine the extent to which the binding partner bound to the FGF-L polypeptide. Typically, a molecule will be tested over a range of concentrations, and a series of control wells lacking one or more elements of the test assays can be used for accuracy in the evaluation of the results. An alternative to this method involves reversing the "positions" of the proteins, *i.e.*, immobilizing FGF-L polypeptide binding partner to the microtiter plate wells, incubating with the test molecule and radiolabeled FGF-L polypeptide, and determining the extent of FGF-L polypeptide binding. *See, e.g., Current Protocols in Molecular Biology*, chap. 18 (Ausubel *et al.*, eds., Green Publishers Inc. and Wiley and Sons 1995).

As an alternative to radiolabeling, an FGF-L polypeptide or its binding partner may be conjugated to biotin, and the presence of biotinylated protein can then be detected using streptavidin linked to an enzyme, such as horse radish peroxidase (HRP) or alkaline phosphatase (AP), which can be detected colorimetrically, or by fluorescent tagging of streptavidin. An antibody directed to an FGF-L polypeptide or to an FGF-L polypeptide binding partner, and which is conjugated to biotin, may also be used for purposes of detection

following incubation of the complex with enzyme-linked streptavidin linked to AP or HRP.

A FGF-L polypeptide or an FGF-L polypeptide binding partner can also be immobilized by attachment to agarose beads, acrylic beads, or other types of such inert solid phase substrates. The substrate-protein complex can be placed in a solution containing the complementary protein and the test compound. After incubation, the beads can be precipitated by centrifugation, and the amount of binding between an FGF-L polypeptide and its binding partner can be assessed using the methods described herein. Alternatively, the substrate-protein complex can be immobilized in a column with the test molecule and complementary protein passing through the column. The formation of a complex between an FGF-L polypeptide and its binding partner can then be assessed using any of the techniques described herein (e.g., radiolabelling or antibody binding).

Another *in vitro* assay that is useful for identifying a test molecule which increases or decreases the formation of a complex between an FGF-L polypeptide binding protein and an FGF-L polypeptide binding partner is a surface plasmon resonance detector system such as the BIAcore assay system (Pharmacia, Piscataway, NJ). The BIAcore system is utilized as specified by the manufacturer. This assay essentially involves the covalent binding of either FGF-L polypeptide or an FGF-L polypeptide binding partner to a dextran-coated sensor chip that is located in a detector. The test compound and the other complementary protein can then be injected, either simultaneously or sequentially, into the chamber containing the sensor chip. The amount of complementary protein that binds can be assessed based on the change in molecular mass that is physically associated with the dextran-coated side of the sensor chip, with the change in molecular mass being measured by the detector system.

In some cases, it may be desirable to evaluate two or more test compounds together for their ability to increase or decrease the formation of a complex between an FGF-L polypeptide and an FGF-L polypeptide binding partner. In these cases, the assays set forth herein can be readily modified by

adding such additional test compound(s) either simultaneously with, or subsequent to, the first test compound. The remainder of the steps in the assay are as set forth herein.

In vitro assays such as those described herein may be used advantageously to screen large numbers of compounds for an effect on the formation of a complex between an FGF-L polypeptide and FGF-L polypeptide binding partner. The assays may be automated to screen compounds generated in phage display, synthetic peptide, and chemical synthesis libraries.

Compounds which increase or decrease the formation of a complex between an FGF-L polypeptide and an FGF-L polypeptide binding partner may also be screened in cell culture using cells and cell lines expressing either FGF-L polypeptide or FGF-L polypeptide binding partner. Cells and cell lines may be obtained from any mammal, but preferably will be from human or other primate, canine, or rodent sources. The binding of an FGF-L polypeptide to cells expressing FGF-L polypeptide binding partner at the surface is evaluated in the presence or absence of test molecules, and the extent of binding may be determined by, for example, flow cytometry using a biotinylated antibody to an FGF-L polypeptide binding partner. Cell culture assays can be used advantageously to further evaluate compounds that score positive in protein binding assays described herein.

Cell cultures can also be used to screen the impact of a drug candidate. For example, drug candidates may decrease or increase the expression of the FGF-L gene. In certain embodiments, the amount of FGF-L polypeptide or an FGF-L polypeptide fragment that is produced may be measured after exposure of the cell culture to the drug candidate. In certain embodiments, one may detect the actual impact of the drug candidate on the cell culture. For example, the over-expression of a particular gene may have a particular impact on the cell culture. In such cases, one may test a drug candidate's ability to increase or decrease the expression of the gene or its ability to prevent or inhibit a particular impact on the cell culture. In other examples, the production of a particular metabolic product such as a fragment of a

polypeptide, may result in, or be associated with, a disease or pathological condition. In such cases, one may test a drug candidate's ability to decrease the production of such a metabolic product in a cell culture.

5 Internalizing Proteins

The *tat* protein sequence (from HIV) can be used to internalize proteins into a cell. See, e.g., Falwell *et al.*, 1994, *Proc. Natl. Acad. Sci. U.S.A.* 91:664-68. For example, an 11 amino acid sequence (Y-G-R-K-K-R-R-Q-R-R-R; SEQ ID NO: 8) of the HIV *tat* protein (termed the "protein
10 transduction domain," or TAT PDT) has been described as mediating delivery across the cytoplasmic membrane and the nuclear membrane of a cell. See Schwarze *et al.*, 1999, *Science* 285:1569-72; and Nagahara *et al.*, 1998, *Nat. Med.* 4:1449-52. In these procedures, FITC-constructs (FITC-labeled G-G-G-G-Y-G-R-K-K-R-R-Q-R-R-R; SEQ ID NO: 9), which penetrate tissues
15 following intraperitoneal administration, are prepared, and the binding of such constructs to cells is detected by fluorescence-activated cell sorting (FACS) analysis. Cells treated with a *tat*- β -gal fusion protein will demonstrate β -gal activity. Following injection, expression of such a construct can be detected in a number of tissues, including liver, kidney, lung, heart, and brain tissue. It is
20 believed that such constructs undergo some degree of unfolding in order to enter the cell, and as such, may require a refolding following entry into the cell.

It will thus be appreciated that the *tat* protein sequence may be used to internalize a desired polypeptide into a cell. For example, using the *tat* protein
25 sequence, an FGF-L antagonist (such as an anti-FGF-L selective binding agent, small molecule, soluble receptor, or antisense oligonucleotide) can be administered intracellularly to inhibit the activity of an FGF-L molecule. As used herein, the term "FGF-L molecule" refers to both FGF-L nucleic acid molecules and FGF-L polypeptides as defined herein. Where desired, the
30 FGF-L protein itself may also be internally administered to a cell using these procedures. See also, Straus, 1999, *Science* 285:1466-67.

Cell Source Identification Using FGF-L Polypeptide

In accordance with certain embodiments of the invention, it may be useful to be able to determine the source of a certain cell type associated with an FGF-L polypeptide. For example, it may be useful to determine the origin
5 of a disease or pathological condition as an aid in selecting an appropriate therapy. In certain embodiments, nucleic acids encoding an FGF-L polypeptide can be used as a probe to identify cells described herein by screening the nucleic acids of the cells with such a probe. In other
10 embodiments, one may use anti-FGF-L polypeptide antibodies to test for the presence of FGF-L polypeptide in cells, and thus, determine if such cells are of the types described herein.

FGF-L Polypeptide Compositions and Administration

Therapeutic compositions are within the scope of the present invention.
15 Such FGF-L polypeptide pharmaceutical compositions may comprise a therapeutically effective amount of an FGF-L polypeptide or an FGF-L nucleic acid molecule in admixture with a pharmaceutically or physiologically acceptable formulation agent selected for suitability with the mode of
20 administration. Pharmaceutical compositions may comprise a therapeutically effective amount of one or more FGF-L polypeptide selective binding agents in admixture with a pharmaceutically or physiologically acceptable formulation agent selected for suitability with the mode of administration.

Acceptable formulation materials preferably are nontoxic to recipients at the dosages and concentrations employed.

25 The pharmaceutical composition may contain formulation materials for modifying, maintaining, or preserving, for example, the pH, osmolarity, viscosity, clarity, color, isotonicity, odor, sterility, stability, rate of dissolution or release, adsorption, or penetration of the composition. Suitable formulation materials include, but are not limited to, amino acids (such as glycine,
30 glutamine, asparagine, arginine, or lysine), antimicrobials, antioxidants (such as ascorbic acid, sodium sulfite, or sodium hydrogen-sulfite), buffers (such as borate, bicarbonate, Tris-HCl, citrates, phosphates, or other organic acids),

bulking agents (such as mannitol or glycine), chelating agents (such as ethylenediamine tetraacetic acid (EDTA)), complexing agents (such as caffeine, polyvinylpyrrolidone, beta-cyclodextrin, or hydroxypropyl-beta-cyclodextrin), fillers, monosaccharides, disaccharides, and other carbohydrates
5 (such as glucose, mannose, or dextrans), proteins (such as serum albumin, gelatin, or immunoglobulins), coloring, flavoring and diluting agents, emulsifying agents, hydrophilic polymers (such as polyvinylpyrrolidone), low molecular weight polypeptides, salt-forming counterions (such as sodium), preservatives (such as benzalkonium FGF-Loride, benzoic acid, salicylic acid,
10 thimerosal, phenethyl alcohol, methylparaben, propylparaben, FGF-Lorhexidine, sorbic acid, or hydrogen peroxide), solvents (such as glycerin, propylene glycol, or polyethylene glycol), sugar alcohols (such as mannitol or sorbitol), suspending agents, surfactants or wetting agents (such as pluronics; PEG; sorbitan esters; polysorbates such as polysorbate 20 or polysorbate 80;
15 triton; tromethamine; lecithin; cholesterol or tyloxapal), stability enhancing agents (such as sucrose or sorbitol), tonicity enhancing agents (such as alkali metal halides – preferably sodium or potassium FGF-Loride – or mannitol sorbitol), delivery vehicles, diluents, excipients and/or pharmaceutical adjuvants. *See Remington's Pharmaceutical Sciences* (18th Ed., A.R. Gennaro, ed., Mack Publishing Company 1990.

The optimal pharmaceutical composition will be determined by a skilled artisan depending upon, for example, the intended route of administration, delivery format, and desired dosage. *See, e.g., Remington's Pharmaceutical Sciences, supra.* Such compositions may influence the
25 physical state, stability, rate of *in vivo* release, and rate of *in vivo* clearance of the FGF-L molecule.

The primary vehicle or carrier in a pharmaceutical composition may be either aqueous or non-aqueous in nature. For example, a suitable vehicle or carrier for injection may be water, physiological saline solution, or artificial
30 cerebrospinal fluid, possibly supplemented with other materials common in compositions for parenteral administration. Neutral buffered saline or saline mixed with serum albumin are further exemplary vehicles. Other exemplary

pharmaceutical compositions comprise Tris buffer of about pH 7.0-8.5, or acetate buffer of about pH 4.0-5.5, which may further include sorbitol or a suitable substitute. In one embodiment of the present invention, FGF-L polypeptide compositions may be prepared for storage by mixing the selected composition having the desired degree of purity with optional formulation agents (*Remington's Pharmaceutical Sciences, supra*) in the form of a lyophilized cake or an aqueous solution. Further, the FGF-L polypeptide product may be formulated as a lyophilizate using appropriate excipients such as sucrose.

10 The FGF-L polypeptide pharmaceutical compositions can be selected for parenteral delivery. Alternatively, the compositions may be selected for inhalation or for delivery through the digestive tract, such as orally. The preparation of such pharmaceutically acceptable compositions is within the skill of the art.

15 The formulation components are present in concentrations that are acceptable to the site of administration. For example, buffers are used to maintain the composition at physiological pH or at a slightly lower pH, typically within a pH range of from about 5 to about 8.

When parenteral administration is contemplated, the therapeutic compositions for use in this invention may be in the form of a pyrogen-free, parenterally acceptable, aqueous solution comprising the desired FGF-L molecule in a pharmaceutically acceptable vehicle. A particularly suitable vehicle for parenteral injection is sterile distilled water in which an FGF-L molecule is formulated as a sterile, isotonic solution, properly preserved. Yet another preparation can involve the formulation of the desired molecule with an agent, such as injectable microspheres, bio-erodible particles, polymeric compounds (such as polylactic acid or polyglycolic acid), beads, or liposomes, that provides for the controlled or sustained release of the product which may then be delivered via a depot injection. Hyaluronic acid may also be used, and this may have the effect of promoting sustained duration in the circulation. Other suitable means for the introduction of the desired molecule include implantable drug delivery devices.

In one embodiment, a pharmaceutical composition may be formulated for inhalation. For example, FGF-L polypeptide may be formulated as a dry powder for inhalation. FGF-L polypeptide or nucleic acid molecule inhalation solutions may also be formulated with a propellant for aerosol delivery. In yet
5 another embodiment, solutions may be nebulized. Pulmonary administration is further described in PCT Pub. No. WO 94/20069, which describes the pulmonary delivery of chemically modified proteins.

It is also contemplated that certain formulations may be administered orally. In one embodiment of the present invention, FGF-L polypeptides that
10 are administered in this fashion can be formulated with or without those carriers customarily used in the compounding of solid dosage forms such as tablets and capsules. For example, a capsule may be designed to release the active portion of the formulation at the point in the gastrointestinal tract when bioavailability is maximized and pre-systemic degradation is minimized.
15 Additional agents can be included to facilitate absorption of the FGF-L polypeptide. Diluents, flavorings, low melting point waxes, vegetable oils, lubricants, suspending agents, tablet disintegrating agents, and binders may also be employed.

Another pharmaceutical composition may involve an effective quantity
20 of FGF-L polypeptides in a mixture with non-toxic excipients that are suitable for the manufacture of tablets. By dissolving the tablets in sterile water, or another appropriate vehicle, solutions can be prepared in unit-dose form. Suitable excipients include, but are not limited to, inert diluents, such as calcium carbonate, sodium carbonate or bicarbonate, lactose, or calcium
25 phosphate; or binding agents, such as starch, gelatin, or acacia; or lubricating agents such as magnesium stearate, stearic acid, or talc.

Additional FGF-L polypeptide pharmaceutical compositions will be evident to those skilled in the art, including formulations involving FGF-L polypeptides in sustained- or controlled-delivery formulations. Techniques for
30 formulating a variety of other sustained- or controlled-delivery means, such as liposome carriers, bio-erodible microparticles or porous beads and depot injections, are also known to those skilled in the art. *See, e.g.,*

PCT/US93/00829, which describes the controlled release of porous polymeric microparticles for the delivery of pharmaceutical compositions.

Additional examples of sustained-release preparations include semipermeable polymer matrices in the form of shaped articles, *e.g.* films, or
5 microcapsules. Sustained release matrices may include polyesters, hydrogels, polylactides (U.S. Patent No. 3,773,919 and European Patent No. 058481), copolymers of L-glutamic acid and gamma ethyl-L-glutamate (Sidman *et al.*, 1983, *Biopolymers* 22:547-56), poly(2-hydroxyethyl-methacrylate) (Langer *et al.*, 1981, *J. Biomed. Mater. Res.* 15:167-277 and Langer, 1982, *Chem. Tech.*
10 12:98-105), ethylene vinyl acetate (Langer *et al.*, *supra*) or poly-D(-)-3-hydroxybutyric acid (European Patent No. 133988). Sustained-release compositions may also include liposomes, which can be prepared by any of several methods known in the art. *See, e.g.*, Eppstein *et al.*, 1985, *Proc. Natl. Acad. Sci. USA* 82:3688-92; and European Patent Nos. 036676, 088046, and
15 143949.

The FGF-L pharmaceutical composition to be used for *in vivo* administration typically must be sterile. This may be accomplished by filtration through sterile filtration membranes. Where the composition is lyophilized, sterilization using this method may be conducted either prior to,
20 or following, lyophilization and reconstitution. The composition for parenteral administration may be stored in lyophilized form or in a solution. In addition, parenteral compositions generally are placed into a container having a sterile access port, for example, an intravenous solution bag or vial having a stopper pierceable by a hypodermic injection needle.

25 Once the pharmaceutical composition has been formulated, it may be stored in sterile vials as a solution, suspension, gel, emulsion, solid, or as a dehydrated or lyophilized powder. Such formulations may be stored either in a ready-to-use form or in a form (*e.g.*, lyophilized) requiring reconstitution prior to administration.

30 In a specific embodiment, the present invention is directed to kits for producing a single-dose administration unit. The kits may each contain both a first container having a dried protein and a second container having an

aqueous formulation. Also included within the scope of this invention are kits containing single and multi-chambered pre-filled syringes (*e.g.*, liquid syringes and lyosyringes).

The effective amount of an FGF-L pharmaceutical composition to be employed therapeutically will depend, for example, upon the therapeutic context and objectives. One skilled in the art will appreciate that the appropriate dosage levels for treatment will thus vary depending, in part, upon the molecule delivered, the indication for which the FGF-L molecule is being used, the route of administration, and the size (body weight, body surface, or organ size) and condition (the age and general health) of the patient. Accordingly, the clinician may titer the dosage and modify the route of administration to obtain the optimal therapeutic effect. A typical dosage may range from about 0.1 $\mu\text{g/kg}$ to up to about 100 mg/kg or more, depending on the factors mentioned above. In other embodiments, the dosage may range from 0.1 $\mu\text{g/kg}$ up to about 100 mg/kg; or 1 $\mu\text{g/kg}$ up to about 100 mg/kg; or 5 $\mu\text{g/kg}$ up to about 100 mg/kg.

The frequency of dosing will depend upon the pharmacokinetic parameters of the FGF-L molecule in the formulation being used. Typically, a clinician will administer the composition until a dosage is reached that achieves the desired effect. The composition may therefore be administered as a single dose, as two or more doses (which may or may not contain the same amount of the desired molecule) over time, or as a continuous infusion via an implantation device or catheter. Further refinement of the appropriate dosage is routinely made by those of ordinary skill in the art and is within the ambit of tasks routinely performed by them. Appropriate dosages may be ascertained through use of appropriate dose-response data.

The route of administration of the pharmaceutical composition is in accord with known methods, *e.g.*, orally; through injection by intravenous, intraperitoneal, intracerebral (intraparenchymal), intracerebroventricular, intramuscular, intraocular, intraarterial, intraportal, or intralesional routes; by sustained release systems; or by implantation devices. Where desired, the

compositions may be administered by bolus injection or continuously by infusion, or by implantation device.

Alternatively or additionally, the composition may be administered locally via implantation of a membrane, sponge, or other appropriate material onto which the desired molecule has been absorbed or encapsulated. Where
5 an implantation device is used, the device may be implanted into any suitable tissue or organ, and delivery of the desired molecule may be via diffusion, timed-release bolus, or continuous administration.

In some cases, it may be desirable to use FGF-L polypeptide pharmaceutical compositions in an *ex vivo* manner. In such instances, cells,
10 tissues, or organs that have been removed from the patient are exposed to FGF-L polypeptide pharmaceutical compositions after which the cells, tissues, or organs are subsequently implanted back into the patient.

In other cases, an FGF-L polypeptide can be delivered by implanting
15 certain cells that have been genetically engineered, using methods such as those described herein, to express and secrete the FGF-L polypeptide. Such cells may be animal or human cells, and may be autologous, heterologous, or xenogeneic. Optionally, the cells may be immortalized. In order to decrease the chance of an immunological response, the cells may be encapsulated to
20 avoid infiltration of surrounding tissues. The encapsulation materials are typically biocompatible, semi-permeable polymeric enclosures or membranes that allow the release of the protein product(s) but prevent the destruction of the cells by the patient's immune system or by other detrimental factors from the surrounding tissues.

As discussed herein, it may be desirable to treat isolated cell
25 populations (such as stem cells, lymphocytes, red blood cells, chondrocytes, neurons, and the like) with one or more FGF-L polypeptides. This can be accomplished by exposing the isolated cells to the polypeptide directly, where it is in a form that is permeable to the cell membrane.

30 Additional embodiments of the present invention relate to cells and methods (*e.g.*, homologous recombination and/or other recombinant production methods) for both the *in vitro* production of therapeutic

polypeptides and for the production and delivery of therapeutic polypeptides by gene therapy or cell therapy. Homologous and other recombination methods may be used to modify a cell that contains a normally transcriptionally-silent FGF-L gene, or an under-expressed gene, and thereby
5 produce a cell which expresses therapeutically efficacious amounts of FGF-L polypeptides.

Homologous recombination is a technique originally developed for targeting genes to induce or correct mutations in transcriptionally active genes. Kucherlapati, 1989, *Prog. in Nucl. Acid Res. & Mol. Biol.* 36:301. The basic
10 technique was developed as a method for introducing specific mutations into specific regions of the mammalian genome (Thomas *et al.*, 1986, *Cell* 44:419-28; Thomas and Capecchi, 1987, *Cell* 51:503-12; Doetschman *et al.*, 1988, *Proc. Natl. Acad. Sci. U.S.A.* 85:8583-87) or to correct specific mutations within defective genes (Doetschman *et al.*, 1987, *Nature* 330:576-78).
15 Exemplary homologous recombination techniques are described in U.S. Patent No. 5,272,071; European Patent Nos. 9193051 and 505500; PCT/US90/07642, and PCT Pub No. WO 91/09955).

Through homologous recombination, the DNA sequence to be inserted into the genome can be directed to a specific region of the gene of interest by
20 attaching it to targeting DNA. The targeting DNA is a nucleotide sequence that is complementary (homologous) to a region of the genomic DNA. Small pieces of targeting DNA that are complementary to a specific region of the genome are put in contact with the parental strand during the DNA replication process. It is a general property of DNA that has been inserted into a cell to
25 hybridize, and therefore, recombine with other pieces of endogenous DNA through shared homologous regions. If this complementary strand is attached to an oligonucleotide that contains a mutation or a different sequence or an additional nucleotide, it too is incorporated into the newly synthesized strand as a result of the recombination. As a result of the proofreading function, it is
30 possible for the new sequence of DNA to serve as the template. Thus, the transferred DNA is incorporated into the genome.

Attached to these pieces of targeting DNA are regions of DNA that may interact with or control the expression of an FGF-L polypeptide, *e.g.*, flanking sequences. For example, a promoter/enhancer element, a suppressor, or an exogenous transcription modulatory element is inserted in the genome of the intended host cell in proximity and orientation sufficient to influence the transcription of DNA encoding the desired FGF-L polypeptide. The control element controls a portion of the DNA present in the host cell genome. Thus, the expression of the desired FGF-L polypeptide may be achieved not by transfection of DNA that encodes the FGF-L gene itself, but rather by the use of targeting DNA (containing regions of homology with the endogenous gene of interest) coupled with DNA regulatory segments that provide the endogenous gene sequence with recognizable signals for transcription of an FGF-L gene.

In an exemplary method, the expression of a desired targeted gene in a cell (*i.e.*, a desired endogenous cellular gene) is altered via homologous recombination into the cellular genome at a preselected site, by the introduction of DNA which includes at least a regulatory sequence, an exon, and a splice donor site. These components are introduced into the chromosomal (genomic) DNA in such a manner that this, in effect, results in the production of a new transcription unit (in which the regulatory sequence, the exon, and the splice donor site present in the DNA construct are operatively linked to the endogenous gene). As a result of the introduction of these components into the chromosomal DNA, the expression of the desired endogenous gene is altered.

Altered gene expression, as described herein, encompasses activating (or causing to be expressed) a gene which is normally silent (unexpressed) in the cell as obtained, as well as increasing the expression of a gene which is not expressed at physiologically significant levels in the cell as obtained. The embodiments further encompass changing the pattern of regulation or induction such that it is different from the pattern of regulation or induction that occurs in the cell as obtained, and reducing (including eliminating) the expression of a gene which is expressed in the cell as obtained.

One method by which homologous recombination can be used to increase, or cause, FGF-L polypeptide production from a cell's endogenous FGF-L gene involves first using homologous recombination to place a recombination sequence from a site-specific recombination system (e.g.,
5 Cre/loxP, FLP/FRT) (Sauer, 1994, *Curr. Opin. Biotechnol.*, 5:521-27; Sauer, 1993, *Methods Enzymol.*, 225:890-900) upstream of (i.e., 5' to) the cell's endogenous genomic FGF-L polypeptide coding region. A plasmid containing a recombination site homologous to the site that was placed just upstream of the genomic FGF-L polypeptide coding region is introduced into the modified
10 cell line along with the appropriate recombinase enzyme. This recombinase causes the plasmid to integrate, via the plasmid's recombination site, into the recombination site located just upstream of the genomic FGF-L polypeptide coding region in the cell line (Baubonis and Sauer, 1993, *Nucleic Acids Res.* 21:2025-29; O'Gorman *et al.*, 1991, *Science* 251:1351-55). Any flanking
15 sequences known to increase transcription (e.g., enhancer/promoter, intron, translational enhancer), if properly positioned in this plasmid, would integrate in such a manner as to create a new or modified transcriptional unit resulting in *de novo* or increased FGF-L polypeptide production from the cell's endogenous FGF-L gene.

20 A further method to use the cell line in which the site specific recombination sequence had been placed just upstream of the cell's endogenous genomic FGF-L polypeptide coding region is to use homologous recombination to introduce a second recombination site elsewhere in the cell line's genome. The appropriate recombinase enzyme is then introduced into
25 the two-recombination-site cell line, causing a recombination event (deletion, inversion, and translocation) (Sauer, 1994, *Curr. Opin. Biotechnol.*, 5:521-27; Sauer, 1993, *Methods Enzymol.*, 225:890-900) that would create a new or modified transcriptional unit resulting in *de novo* or increased FGF-L polypeptide production from the cell's endogenous FGF-L gene.

30 An additional approach for increasing, or causing, the expression of FGF-L polypeptide from a cell's endogenous FGF-L gene involves increasing, or causing, the expression of a gene or genes (e.g., transcription factors) and/or

decreasing the expression of a gene or genes (*e.g.*, transcriptional repressors) in a manner which results in *de novo* or increased FGF-L polypeptide production from the cell's endogenous FGF-L gene. This method includes the introduction of a non-naturally occurring polypeptide (*e.g.*, a polypeptide
5 comprising a site specific DNA binding domain fused to a transcriptional factor domain) into the cell such that *de novo* or increased FGF-L polypeptide production from the cell's endogenous FGF-L gene results.

The present invention further relates to DNA constructs useful in the method of altering expression of a target gene. In certain embodiments, the
10 exemplary DNA constructs comprise: (a) one or more targeting sequences, (b) a regulatory sequence, (c) an exon, and (d) an unpaired splice-donor site. The targeting sequence in the DNA construct directs the integration of elements (a) - (d) into a target gene in a cell such that the elements (b) - (d) are operatively linked to sequences of the endogenous target gene. In another embodiment,
15 the DNA constructs comprise: (a) one or more targeting sequences, (b) a regulatory sequence, (c) an exon, (d) a splice-donor site, (e) an intron, and (f) a splice-acceptor site, wherein the targeting sequence directs the integration of elements (a) - (f) such that the elements of (b) - (f) are operatively linked to the endogenous gene. The targeting sequence is homologous to the preselected
20 site in the cellular chromosomal DNA with which homologous recombination is to occur. In the construct, the exon is generally 3' of the regulatory sequence and the splice-donor site is 3' of the exon.

If the sequence of a particular gene is known, such as the nucleic acid sequence of FGF-L polypeptide presented herein, a piece of DNA that is
25 complementary to a selected region of the gene can be synthesized or otherwise obtained, such as by appropriate restriction of the native DNA at specific recognition sites bounding the region of interest. This piece serves as a targeting sequence upon insertion into the cell and will hybridize to its homologous region within the genome. If this hybridization occurs during
30 DNA replication, this piece of DNA, and any additional sequence attached thereto, will act as an Okazaki fragment and will be incorporated into the newly synthesized daughter strand of DNA. The present invention, therefore,

includes nucleotides encoding an FGF-L polypeptide, which nucleotides may be used as targeting sequences.

5 FGF-L polypeptide cell therapy, *e.g.*, the implantation of cells producing FGF-L polypeptides, is also contemplated. This embodiment involves implanting cells capable of synthesizing and secreting a biologically active form of FGF-L polypeptide. Such FGF-L polypeptide-producing cells can be cells that are natural producers of FGF-L polypeptides or may be recombinant cells whose ability to produce FGF-L polypeptides has been augmented by transformation with a gene encoding the desired FGF-L polypeptide or with a gene augmenting the expression of FGF-L polypeptide. 10 Such a modification may be accomplished by means of a vector suitable for delivering the gene as well as promoting its expression and secretion. In order to minimize a potential immunological reaction in patients being administered an FGF-L polypeptide, as may occur with the administration of a polypeptide of a foreign species, it is preferred that the natural cells producing FGF-L polypeptide be of human origin and produce human FGF-L polypeptide. 15 Likewise, it is preferred that the recombinant cells producing FGF-L polypeptide be transformed with an expression vector containing a gene encoding a human FGF-L polypeptide.

20 Implanted cells may be encapsulated to avoid the infiltration of surrounding tissue. Human or non-human animal cells may be implanted in patients in biocompatible, semipermeable polymeric enclosures or membranes that allow the release of FGF-L polypeptide, but that prevent the destruction of the cells by the patient's immune system or by other detrimental factors from the surrounding tissue. 25 Alternatively, the patient's own cells, transformed to produce FGF-L polypeptides *ex vivo*, may be implanted directly into the patient without such encapsulation.

Techniques for the encapsulation of living cells are known in the art, and the preparation of the encapsulated cells and their implantation in patients 30 may be routinely accomplished. For example, Baetge *et al.* (PCT Pub. No. WO 95/05452 and PCT/US94/09299) describe membrane capsules containing genetically engineered cells for the effective delivery of biologically active

molecules. The capsules are biocompatible and are easily retrievable. The capsules encapsulate cells transfected with recombinant DNA molecules comprising DNA sequences coding for biologically active molecules operatively linked to promoters that are not subject to down-regulation *in vivo* upon implantation into a mammalian host. The devices provide for the delivery of the molecules from living cells to specific sites within a recipient. In addition, see U.S. Patent Nos. 4,892,538; 5,011,472; and 5,106,627. A system for encapsulating living cells is described in PCT Pub. No. WO 91/10425 (Aebischer *et al.*). See also, PCT Pub. No. WO 91/10470 (Aebischer *et al.*); Winn *et al.*, 1991, *Exper. Neurol.* 113:322-29; Aebischer *et al.*, 1991, *Exper. Neurol.* 111:269-75; and Tresco *et al.*, 1992, *ASAIO* 38:17-23.

In vivo and *in vitro* gene therapy delivery of FGF-L polypeptides is also envisioned. One example of a gene therapy technique is to use the FGF-L gene (either genomic DNA, cDNA, and/or synthetic DNA) encoding an FGF-L polypeptide which may be operably linked to a constitutive or inducible promoter to form a "gene therapy DNA construct." The promoter may be homologous or heterologous to the endogenous FGF-L gene, provided that it is active in the cell or tissue type into which the construct will be inserted. Other components of the gene therapy DNA construct may optionally include DNA molecules designed for site-specific integration (*e.g.*, endogenous sequences useful for homologous recombination), tissue-specific promoters, enhancers or silencers, DNA molecules capable of providing a selective advantage over the parent cell, DNA molecules useful as labels to identify transformed cells, negative selection systems, cell specific binding agents (as, for example, for cell targeting), cell-specific internalization factors, transcription factors enhancing expression from a vector, and factors enabling vector production.

A gene therapy DNA construct can then be introduced into cells (either *ex vivo* or *in vivo*) using viral or non-viral vectors. One means for introducing the gene therapy DNA construct is by means of viral vectors as described herein. Certain vectors, such as retroviral vectors, will deliver the DNA construct to the chromosomal DNA of the cells, and the gene can integrate

into the chromosomal DNA. Other vectors will function as episomes, and the gene therapy DNA construct will remain in the cytoplasm.

In yet other embodiments, regulatory elements can be included for the controlled expression of the FGF-L gene in the target cell. Such elements are
5 turned on in response to an appropriate effector. In this way, a therapeutic polypeptide can be expressed when desired. One conventional control means involves the use of small molecule dimerizers or rapalogs to dimerize chimeric proteins which contain a small molecule-binding domain and a domain capable of initiating a biological process, such as a DNA-binding protein or
10 transcriptional activation protein (*see* PCT Pub. Nos. WO 96/41865, WO 97/31898, and WO 97/31899). The dimerization of the proteins can be used to initiate transcription of the transgene.

An alternative regulation technology uses a method of storing proteins expressed from the gene of interest inside the cell as an aggregate or cluster.
15 The gene of interest is expressed as a fusion protein that includes a conditional aggregation domain that results in the retention of the aggregated protein in the endoplasmic reticulum. The stored proteins are stable and inactive inside the cell. The proteins can be released, however, by administering a drug (*e.g.*, small molecule ligand) that removes the conditional aggregation domain and
20 thereby specifically breaks apart the aggregates or clusters so that the proteins may be secreted from the cell. *See* Aridor *et al.*, 2000, *Science* 287:816-17 and Rivera *et al.*, 2000, *Science* 287:826-30.

Other suitable control means or gene switches include, but are not limited to, the systems described herein. Mifepristone (RU486) is used as a
25 progesterone antagonist. The binding of a modified progesterone receptor ligand-binding domain to the progesterone antagonist activates transcription by forming a dimer of two transcription factors that then pass into the nucleus to bind DNA. The ligand-binding domain is modified to eliminate the ability of the receptor to bind to the natural ligand. The modified steroid hormone
30 receptor system is further described in U.S. Patent No. 5,364,791 and PCT Pub. Nos. WO 96/40911 and WO 97/10337.

Yet another control system uses ecdysone (a fruit fly steroid hormone) which binds to and activates an ecdysone receptor (cytoplasmic receptor). The receptor then translocates to the nucleus to bind a specific DNA response element (promoter from ecdysone-responsive gene). The ecdysone receptor
5 includes a transactivation domain, DNA-binding domain, and ligand-binding domain to initiate transcription. The ecdysone system is further described in U.S. Patent No. 5,514,578 and PCT Pub. Nos. WO 97/38117, WO 96/37609, and WO 93/03162.

Another control means uses a positive tetracycline-controllable
10 transactivator. This system involves a mutated tet repressor protein DNA-binding domain (mutated tet R-4 amino acid changes which resulted in a reverse tetracycline-regulated transactivator protein, *i.e.*, it binds to a tet operator in the presence of tetracycline) linked to a polypeptide which activates transcription. Such systems are described in U.S. Patent Nos.
15 5,464,758, 5,650,298, and 5,654,168.

Additional expression control systems and nucleic acid constructs are described in U.S. Patent Nos. 5,741,679 and 5,834,186, to Innovir Laboratories Inc.

In vivo gene therapy may be accomplished by introducing the gene
20 encoding FGF-L polypeptide into cells via local injection of an FGF-L nucleic acid molecule or by other appropriate viral or non-viral delivery vectors. Hefti 1994, *Neurobiology* 25:1418-35. For example, a nucleic acid molecule encoding an FGF-L polypeptide may be contained in an adeno-associated virus (AAV) vector for delivery to the targeted cells (*see, e.g.*, Johnson, PCT
25 Pub. No. WO 95/34670; PCT App. No. PCT/US95/07178). The recombinant AAV genome typically contains AAV inverted terminal repeats flanking a DNA sequence encoding an FGF-L polypeptide operably linked to functional promoter and polyadenylation sequences.

Alternative suitable viral vectors include, but are not limited to,
30 retrovirus, adenovirus, herpes simplex virus, lentivirus, hepatitis virus, parvovirus, papovavirus, poxvirus, alphavirus, coronavirus, rhabdovirus, paramyxovirus, and papilloma virus vectors. U.S. Patent No. 5,672,344

describes an *in vivo* viral-mediated gene transfer system involving a recombinant neurotrophic HSV-1 vector. U.S. Patent No. 5,399,346 provides examples of a process for providing a patient with a therapeutic protein by the delivery of human cells which have been treated *in vitro* to insert a DNA
5 segment encoding a therapeutic protein. Additional methods and materials for the practice of gene therapy techniques are described in U.S. Patent Nos. 5,631,236 (involving adenoviral vectors), 5,672,510 (involving retroviral vectors), 5,635,399 (involving retroviral vectors expressing cytokines).

Nonviral delivery methods include, but are not limited to, liposome-
10 mediated transfer, naked DNA delivery (direct injection), receptor-mediated transfer (ligand-DNA complex), electroporation, calcium phosphate precipitation, and microparticle bombardment (*e.g.*, gene gun). Gene therapy materials and methods may also include inducible promoters, tissue-specific enhancer-promoters, DNA sequences designed for site-specific integration,
15 DNA sequences capable of providing a selective advantage over the parent cell, labels to identify transformed cells, negative selection systems and expression control systems (safety measures), cell-specific binding agents (for cell targeting), cell-specific internalization factors, and transcription factors to enhance expression by a vector as well as methods of vector manufacture.
20 Such additional methods and materials for the practice of gene therapy techniques are described in U.S. Patent Nos. 4,970,154 (involving electroporation techniques), 5,679,559 (describing a lipoprotein-containing system for gene delivery), 5,676,954 (involving liposome carriers), 5,593,875 (describing methods for calcium phosphate transfection), and 4,945,050
25 (describing a process wherein biologically active particles are propelled at cells at a speed whereby the particles penetrate the surface of the cells and become incorporated into the interior of the cells), and PCT Pub. No. WO 96/40958 (involving nuclear ligands).

It is also contemplated that FGF-L gene therapy or cell therapy can
30 further include the delivery of one or more additional polypeptide(s) in the same or a different cell(s). Such cells may be separately introduced into the patient, or the cells may be contained in a single implantable device, such as

the encapsulating membrane described above, or the cells may be separately modified by means of viral vectors.

A means to increase endogenous FGF-L polypeptide expression in a cell via gene therapy is to insert one or more enhancer elements into the FGF-L polypeptide promoter, where the enhancer elements can serve to increase transcriptional activity of the FGF-L gene. The enhancer elements used will be selected based on the tissue in which one desires to activate the gene – enhancer elements known to confer promoter activation in that tissue will be selected. For example, if a gene encoding an FGF-L polypeptide is to be “turned on” in T-cells, the *lck* promoter enhancer element may be used. Here, the functional portion of the transcriptional element to be added may be inserted into a fragment of DNA containing the FGF-L polypeptide promoter (and optionally, inserted into a vector and/or 5’ and/or 3’ flanking sequences) using standard cloning techniques. This construct, known as a “homologous recombination construct,” can then be introduced into the desired cells either *ex vivo* or *in vivo*.

Gene therapy also can be used to decrease FGF-L polypeptide expression by modifying the nucleotide sequence of the endogenous promoter. Such modification is typically accomplished via homologous recombination methods. For example, a DNA molecule containing all or a portion of the promoter of the FGF-L gene selected for inactivation can be engineered to remove and/or replace pieces of the promoter that regulate transcription. For example, the TATA box and/or the binding site of a transcriptional activator of the promoter may be deleted using standard molecular biology techniques; such deletion can inhibit promoter activity thereby repressing the transcription of the corresponding FGF-L gene. The deletion of the TATA box or the transcription activator binding site in the promoter may be accomplished by generating a DNA construct comprising all or the relevant portion of the FGF-L polypeptide promoter (from the same or a related species as the FGF-L gene to be regulated) in which one or more of the TATA box and/or transcriptional activator binding site nucleotides are mutated via substitution, deletion and/or insertion of one or more nucleotides. As a result, the TATA box and/or

activator binding site has decreased activity or is rendered completely inactive. This construct, which also will typically contain at least about 500 bases of DNA that correspond to the native (endogenous) 5' and 3' DNA sequences adjacent to the promoter segment that has been modified, may be
5 introduced into the appropriate cells (either *ex vivo* or *in vivo*) either directly or via a viral vector as described herein. Typically, the integration of the construct into the genomic DNA of the cells will be via homologous recombination, where the 5' and 3' DNA sequences in the promoter construct can serve to help integrate the modified promoter region via hybridization to
10 the endogenous chromosomal DNA.

Therapeutic Uses

FGF-L nucleic acid molecules, polypeptides, and agonists and antagonists thereof can be used to treat, diagnose, ameliorate, or prevent a
15 number of diseases, disorders, or conditions, including those recited herein.

FGF-L polypeptide agonists and antagonists include those molecules which regulate FGF-L polypeptide activity and either increase or decrease at least one activity of the mature form of the FGF-L polypeptide. Agonists or antagonists may be co-factors, such as a protein, peptide, carbohydrate, lipid,
20 or small molecular weight molecule, which interact with FGF-L polypeptide and thereby regulate its activity. Potential polypeptide agonists or antagonists include antibodies that react with either soluble or membrane-bound forms of FGF-L polypeptides that comprise part or all of the extracellular domains of the said proteins. Molecules that regulate FGF-L polypeptide expression
25 typically include nucleic acids encoding FGF-L polypeptide that can act as anti-sense regulators of expression.

The FGF-L nucleic acid molecules, polypeptides, and agonists and antagonists thereof of the present invention are useful for the same purposes for which members of the FGF family of polypeptides are known to be useful.
30 Thus, the FGF-L polypeptides of this invention are potent mitogens for a variety of cells of the mesodermal, ectodermal, and endodermal origin, including fibroblasts, corneal and vascular endothelial cells, granulocytes,

adrenal cortical cells, chondrocytes, myoblasts, vascular smooth muscle cells, lens epithelial cells, melanocytes, keratinocytes, oligodendrocytes, astrocytes, osteoblasts, and hematopoietic cells. Included among these biological activities are the ability to stimulate the proliferation of vascular endothelial cells and to enable endothelial cells to penetrate the basement membrane. Consistent with these properties, the FGF-L polypeptides of this invention may stimulate angiogenesis and promote wound healing (*i.e.*, facilitate the repair or replacement of damages of diseased tissue resulting from burns, traumatic injuries, surgery, or ulcers). These polypeptides may also induce mesoderm formation and modulate the differentiation of neuronal cells, adipocytes, and skeletal muscle cells. The polypeptides may also be employed to prevent or ameliorate skin aging due to sun exposure by stimulating keratinocyte growth. Further, the polypeptides of this invention may be employed to maintain organs before transplantation or for supporting cultures of primary cells and tissues. In addition, these polypeptides may be utilized to prevent hair loss since FGF family members activate hair-forming cells and promote melanocyte growth. They may also be used to stimulate the growth and differentiation of hematopoietic cells and bone marrow cells when used in combination with other cytokines.

The FGF-L gene appears to be most closely related to the FGF-9 and FGF-16 genes. FGF-9, which was initially referred to as glia-activating factor (Miyamoto *et al.*, 1993, *Mol. Cell. Biol.* 13:4251-59), has been localized to the cells of the central nervous system, suggesting that this member of the FGF family plays a role in the central nervous system (Tagashira *et al.*, 1995, *Brain Res. Mol. Brain Res.* 30:233-41). Specifically, it has been suggested that FGF-9 may be involved in the neuropathology of Alzheimer's disease (Nakamura *et al.*, 1998, *Brain Res.* 814:222-25) and the pathogenesis of motor neuron disorders in the adults, especially amyotrophic lateral sclerosis (Kanda *et al.*, 1999, *Int. J. Dev. Neurosci.* 17:191-200). FGF-9 may also play a role in the stimulation of thrombopoiesis (Matsumoto-Yoshitomi *et al.*, 1995, *Growth Factors* 12:179-90) and the odontogenesis (Kettunen and Thesleff, 1998, *Dev. Dyn.* 211:256-68). FGF-16, which has been shown to be expressed in adult

heart and embryonic brown adipose tissue (Miyake *et al.*, 1998, *Biochem. Biophys. Res. Commun.* 243:148-52), may play a role in oligodendrocyte and hepatocellular proliferation (Danilenko *et al.*, 1999, *Arch. Biochem. Biophys.* 361:34-46). Accordingly, FGF-L nucleic acid molecules, polypeptides, and agonists and antagonists thereof may be used to treat, diagnose, ameliorate, or prevent diseases, disorders, or conditions involving the central nervous system, teeth, brown adipose tissue, heart, or liver.

A non-exclusive list of other diseases, disorders, or conditions which may be treated, diagnosed, ameliorated, or prevented with the FGF-L nucleic acid molecules, polypeptides, and agonists and antagonists thereof of the present invention include: dermal wounds, epidermalysis bullosa, male pattern alopecia, gastric ulcer, duodenal ulcer, erosive gastritis, esophagitis, esophageal reflux disease, inflammatory bowel disease, radiation- or chemotherapy-induced gut toxicity, hyaline membrane disease, necrosis of the respiratory epithelium, emphysema, pulmonary inflammation, pulmonary fibrosis, hepatic cirrhosis, fulminant liver failure, viral hepatitis, and diabetes.

Agonists or antagonists of FGF-L polypeptide function may be used (simultaneously or sequentially) in combination with one or more cytokines, growth factors, antibiotics, anti-inflammatories, and/or chemotherapeutic agents as is appropriate for the condition being treated.

Other diseases caused by or mediated by undesirable levels of FGF-L polypeptides are encompassed within the scope of the invention. Undesirable levels include excessive levels of FGF-L polypeptides and sub-normal levels of FGF-L polypeptides.

Uses of FGF-L Nucleic Acids and Polypeptides

Nucleic acid molecules of the invention (including those that do not themselves encode biologically active polypeptides) may be used to map the locations of the FGF-L gene and related genes on chromosomes. Mapping may be done by techniques known in the art, such as PCR amplification and *in situ* hybridization.

FGF-L nucleic acid molecules (including those that do not themselves

encode biologically active polypeptides), may be useful as hybridization probes in diagnostic assays to test, either qualitatively or quantitatively, for the presence of an FGF-L nucleic acid molecule in mammalian tissue or bodily fluid samples.

5 Other methods may also be employed where it is desirable to inhibit the activity of one or more FGF-L polypeptides. Such inhibition may be effected by nucleic acid molecules that are complementary to and hybridize to expression control sequences (triple helix formation) or to FGF-L mRNA. For example, antisense DNA or RNA molecules, which have a sequence that is
10 complementary to at least a portion of an FGF-L gene can be introduced into the cell. Anti-sense probes may be designed by available techniques using the sequence of the FGF-L gene disclosed herein. Typically, each such antisense molecule will be complementary to the start site (5' end) of each selected FGF-L gene. When the antisense molecule then hybridizes to the
15 corresponding FGF-L mRNA, translation of this mRNA is prevented or reduced. Anti-sense inhibitors provide information relating to the decrease or absence of an FGF-L polypeptide in a cell or organism.

 Alternatively, gene therapy may be employed to create a dominant-negative inhibitor of one or more FGF-L polypeptides. In this situation, the
20 DNA encoding a mutant polypeptide of each selected FGF-L polypeptide can be prepared and introduced into the cells of a patient using either viral or non-viral methods as described herein. Each such mutant is typically designed to compete with endogenous polypeptide in its biological role.

 In addition, an FGF-L polypeptide, whether biologically active or not,
25 may be used as an immunogen, that is, the polypeptide contains at least one epitope to which antibodies may be raised. Selective binding agents that bind to an FGF-L polypeptide (as described herein) may be used for *in vivo* and *in vitro* diagnostic purposes, including, but not limited to, use in labeled form to detect the presence of FGF-L polypeptide in a body fluid or cell sample. The
30 antibodies may also be used to prevent, treat, or diagnose a number of diseases and disorders, including those recited herein. The antibodies may bind to an FGF-L polypeptide so as to diminish or block at least one activity

characteristic of an FGF-L polypeptide, or may bind to a polypeptide to increase at least one activity characteristic of an FGF-L polypeptide (including by increasing the pharmacokinetics of the FGF-L polypeptide).

The FGF-L polypeptides of the present invention can be used to clone
5 FGF-L polypeptide receptors, using an expression cloning strategy. Radiolabeled (¹²⁵Iodine) FGF-L polypeptide or affinity/activity-tagged FGF-L polypeptide (such as an Fc fusion or an alkaline phosphatase fusion) can be used in binding assays to identify a cell type or cell line or tissue that expresses FGF-L polypeptide receptors. RNA isolated from such cells or
10 tissues can be converted to cDNA, cloned into a mammalian expression vector, and transfected into mammalian cells (such as COS or 293 cells) to create an expression library. A radiolabeled or tagged FGF-L polypeptide can then be used as an affinity ligand to identify and isolate from this library the subset of cells that express the FGF-L polypeptide receptors on their surface.
15 DNA can then be isolated from these cells and transfected into mammalian cells to create a secondary expression library in which the fraction of cells expressing FGF-L polypeptide receptors is many-fold higher than in the original library. This enrichment process can be repeated iteratively until a single recombinant clone containing an FGF-L polypeptide receptor is
20 isolated. Isolation of the FGF-L polypeptide receptors is useful for identifying or developing novel agonists and antagonists of the FGF-L polypeptide signaling pathway. Such agonists and antagonists include soluble FGF-L polypeptide receptors, anti-FGF-L polypeptide receptor antibodies, small molecules, or antisense oligonucleotides, and they may be used for treating,
25 preventing, or diagnosing one or more of the diseases or disorders described herein.

The murine and human FGF-L nucleic acids of the present invention are also useful tools for isolating the corresponding chromosomal FGF-L polypeptide genes. For example, mouse chromosomal DNA containing FGF-L
30 L sequences can be used to construct knockout mice, thereby permitting an examination of the *in vivo* role for FGF-L polypeptide. The human FGF-L genomic DNA can be used to identify heritable tissue-degenerating diseases.

A deposit of cDNA encoding human FGF-L polypeptide, subcloned into pGEM-T: JH3 (Promega, Madison, WI), and having Accession No. PTA-1882, was made with the American Type Culture Collection, 10801 University Boulevard, Manassas, VA 20110-2209 on May 17, 2000.

5 The following examples are intended for illustration purposes only, and should not be construed as limiting the scope of the invention in any way.

Example 1: Cloning of the Human FGF-L Polypeptide Gene

Generally, materials and methods as described in Sambrook *et al.*
10 *supra* were used to clone and analyze the gene encoding human FGF-L polypeptide.

A virtual transcript identified in a proprietary genomic database was used to isolate cDNA sequences encoding human FGF-L polypeptide. This virtual transcript was used to design gene specific oligonucleotides for the
15 identification of cDNA sources and the generation of cDNA clones, using various PCR strategies.

The full-length cDNA sequence encoding human FGF-L polypeptide was obtained in amplification reactions using a human fetal stomach cDNA library template and the amplimers 2440-39 (5'-C-C-A-T-G-G-C-T-C-C-C-T-
20 T-A-G-C-C-G-A-A-G-T-C-3'; SEQ ID NO: 10) and 2432-77 (5'-T-C-A-C-T-A-T-C-G-C-A-C-T-T-C-A-A-G-T-G-T-A-C-A-T-C-3'; SEQ ID NO: 11). Reactions were performed at 94°C for 2 minutes for one cycle; 94°C for 45 seconds, 64°C for 45 seconds, and 72°C for 2 minutes for 40 cycles; and 72°C for 7 minutes for one cycle. The amplification mixture was separated on a 1%
25 TBE agarose gel, and a PCR product of approximately 600 bp was isolated from the gel and purified using the Qiagen gel extraction kit. The isolated fragment was subcloned into pGEM-T.

Sequence analysis of the predicted cDNA sequence for human FGF-L polypeptide indicated that the gene comprises a 633 bp open reading frame
30 encoding a protein of 211 amino acids (Figures 1A-1B). Figures 2 and 3 illustrate amino acid sequence alignments of human FGF-L polypeptide and either human or rat FGF-9. Figure 4 illustrates the amino acid sequence

alignment of human FGF-9, human FGF-16, human FGF-L polypeptide, murine FGF-9, and rat FGF-16.

Example 2: FGF-L mRNA Expression

5 The expression of FGF-L mRNA is examined by Northern blot analysis. Multiple human tissue northern blots (Clontech) are probed with a suitable restriction fragment isolated from a human FGF-L polypeptide cDNA clone. The probe is labeled with ^{32}P -dCTP using standard techniques.

10 Northern blots are prehybridized for 2 hours at 42°C in hybridization solution (5X SSC, 50% deionized formamide, 5X Denhardt's solution, 0.5% SDS, and 100 mg/ml denatured salmon sperm DNA) and then hybridized at 42°C overnight in fresh hybridization solution containing 5 ng/ml of the labeled probe. Following hybridization, the filters are washed twice for 10 minutes at room temperature in 2X SSC and 0.1% SDS, and then twice for 30
15 minutes at 65°C in 0.1X SSC and 0.1% SDS. The blots are then exposed to autoradiography.

 The expression of FGF-L mRNA is localized by *in situ* hybridization. A panel of normal embryonic and adult mouse tissues is fixed in 4%
20 paraformaldehyde, embedded in paraffin, and sectioned at 5 μm . Sectioned tissues are permeabilized in 0.2 M HCl, digested with Proteinase K, and acetylated with triethanolamine and acetic anhydride. Sections are prehybridized for 1 hour at 60°C in hybridization solution (300 mM NaCl, 20 mM Tris-HCl, pH 8.0, 5 mM EDTA, 1X Denhardt's solution, 0.2% SDS, 10
25 mM DTT, 0.25 mg/ml tRNA, 25 $\mu\text{g/ml}$ polyA, 25 $\mu\text{g/ml}$ polyC and 50% formamide) and then hybridized overnight at 60°C in the same solution containing 10% dextran and 2×10^4 cpm/ μl of a ^{33}P -labeled antisense riboprobe complementary to the human FGF-L gene. The riboprobe is obtained by *in vitro* transcription of a clone containing human FGF-L cDNA
30 sequences using standard techniques.

 Following hybridization, sections are rinsed in hybridization solution, treated with RNaseA to digest unhybridized probe, and then washed in 0.1X

SSC at 55°C for 30 minutes. Sections are then immersed in NTB-2 emulsion (Kodak, Rochester, NY), exposed for 3 weeks at 4°C, developed, and counterstained with hematoxylin and eosin. Tissue morphology and hybridization signal are simultaneously analyzed by darkfield and standard illumination for brain (one sagittal and two coronal sections), gastrointestinal tract (esophagus, stomach, duodenum, jejunum, ileum, proximal colon, and distal colon), pituitary, liver, lung, heart, spleen, thymus, lymph nodes, kidney, adrenal, bladder, pancreas, salivary gland, male and female reproductive organs (ovary, oviduct, and uterus in the female; and testis, epididymus, prostate, seminal vesicle, and vas deferens in the male), BAT and WAT (subcutaneous, peri-renal), bone (femur), skin, breast, and skeletal muscle.

Example 3: Production of FGF-L Polypeptides

A. Expression of FGF-L Polypeptides in Bacteria

PCR is used to amplify template DNA sequences encoding an FGF-L polypeptide using primers corresponding to the 5' and 3' ends of the sequence. The amplified DNA products may be modified to contain restriction enzyme sites to allow for insertion into expression vectors. PCR products are gel purified and inserted into expression vectors using standard recombinant DNA methodology. An exemplary vector, such as pAMG21 (ATCC no. 98113) containing the lux promoter and a gene encoding kanamycin resistance is digested with Bam HI and Nde I for directional cloning of inserted DNA. The ligated mixture is transformed into an *E. coli* host strain by electroporation and transformants are selected for kanamycin resistance. Plasmid DNA from selected colonies is isolated and subjected to DNA sequencing to confirm the presence of the insert.

Transformed host cells are incubated in 2xYT medium containing 30 µg/mL kanamycin at 30°C prior to induction. Gene expression is induced by the addition of N-(3-oxohexanoyl)-dl-homoserine lactone to a final concentration of 30 ng/mL followed by incubation at either 30°C or 37°C for six hours. The expression of FGF-L polypeptide is evaluated by

centrifugation of the culture, resuspension and lysis of the bacterial pellets, and analysis of host cell proteins by SDS-polyacrylamide gel electrophoresis.

Inclusion bodies containing FGF-L polypeptide are purified as follows. Bacterial cells are pelleted by centrifugation and resuspended in water. The cell suspension is lysed by sonication and pelleted by centrifugation at 195,000 xg for 5 to 10 minutes. The supernatant is discarded, and the pellet is washed and transferred to a homogenizer. The pellet is homogenized in 5 mL of a Percoll solution (75% liquid Percoll and 0.15 M NaCl) until uniformly suspended and then diluted and centrifuged at 21,600 xg for 30 minutes. Gradient fractions containing the inclusion bodies are recovered and pooled. The isolated inclusion bodies are analyzed by SDS-PAGE.

A single band on an SDS polyacrylamide gel corresponding to *E. coli*-produced FGF-L polypeptide is excised from the gel, and the N-terminal amino acid sequence is determined essentially as described by Matsudaira *et al.*, 1987, *J. Biol. Chem.* 262:10-35.

B. Expression of FGF-L Polypeptide in Mammalian Cells

PCR is used to amplify template DNA sequences encoding an FGF-L polypeptide using primers corresponding to the 5' and 3' ends of the sequence. The amplified DNA products may be modified to contain restriction enzyme sites to allow for insertion into expression vectors. PCR products are gel purified and inserted into expression vectors using standard recombinant DNA methodology. An exemplary expression vector, pCEP4 (Invitrogen, Carlsbad, CA), that contains an Epstein-Barr virus origin of replication, may be used for the expression of FGF-L polypeptides in 293-EBNA-1 cells. Amplified and gel purified PCR products are ligated into pCEP4 vector and introduced into 293-EBNA cells by lipofection. The transfected cells are selected in 100 µg/mL hygromycin and the resulting drug-resistant cultures are grown to confluence. The cells are then cultured in serum-free media for 72 hours. The conditioned media is removed and FGF-L polypeptide expression is analyzed by SDS-PAGE.

FGF-L polypeptide expression may be detected by silver staining. Alternatively, FGF-L polypeptide is produced as a fusion protein with an epitope tag, such as an IgG constant domain or a FLAG epitope, which may be detected by Western blot analysis using antibodies to the peptide tag.

5 FGF-L polypeptides may be excised from an SDS-polyacrylamide gel, or FGF-L fusion proteins are purified by affinity chromatography to the epitope tag, and subjected to N-terminal amino acid sequence analysis as described herein.

10 C. Expression and Purification of FGF-L Polypeptide in Mammalian Cells

FGF-L polypeptide expression constructs are introduced into 293 EBNA or CHO cells using either a lipofection or calcium phosphate protocol.

To conduct functional studies on the FGF-L polypeptides that are produced, large quantities of conditioned media are generated from a pool of
15 hygromycin selected 293 EBNA clones. The cells are cultured in 500 cm Nunc Triple Flasks to 80% confluence before switching to serum free media a week prior to harvesting the media. Conditioned media is harvested and frozen at
-20°C until purification.

20 Conditioned media is purified by affinity chromatography as described below. The media is thawed and then passed through a 0.2 µm filter. A Protein G column is equilibrated with PBS at pH 7.0, and then loaded with the filtered media. The column is washed with PBS until the absorbance at A₂₈₀ reaches a baseline. FGF-L polypeptide is eluted from the column with 0.1 M
25 Glycine-HCl at pH 2.7 and immediately neutralized with 1 M Tris-HCl at pH 8.5. Fractions containing FGF-L polypeptide are pooled, dialyzed in PBS, and stored at -70°C.

For Factor Xa cleavage of the human FGF-L polypeptide-Fc fusion polypeptide, affinity chromatography-purified protein is dialyzed in 50 mM
30 Tris-HCl, 100 mM NaCl, 2 mM CaCl₂ at pH 8.0. The restriction protease Factor Xa is added to the dialyzed protein at 1/100 (w/w) and the sample digested overnight at room temperature.

Example 4: Production of Anti-FGF-L Polypeptide Antibodies

Antibodies to FGF-L polypeptides may be obtained by immunization with purified protein or with FGF-L peptides produced by biological or chemical synthesis. Suitable procedures for generating antibodies include those described in Hudson and Bay, *Practical Immunology* (2nd ed., Blackwell Scientific Publications).

In one procedure for the production of antibodies, animals (typically mice or rabbits) are injected with an FGF-L antigen (such as an FGF-L polypeptide), and those with sufficient serum titer levels as determined by ELISA are selected for hybridoma production. Spleens of immunized animals are collected and prepared as single cell suspensions from which splenocytes are recovered. The splenocytes are fused to mouse myeloma cells (such as Sp2/0-Ag14 cells), are first incubated in DMEM with 200 U/mL penicillin, 200 µg/mL streptomycin sulfate, and 4 mM glutamine, and are then incubated in HAT selection medium (hypoxanthine, aminopterin, and thymidine). After selection, the tissue culture supernatants are taken from each fusion well and tested for anti-FGF-L antibody production by ELISA.

Alternative procedures for obtaining anti-FGF-L antibodies may also be employed, such as the immunization of transgenic mice harboring human Ig loci for production of human antibodies, and the screening of synthetic antibody libraries, such as those generated by mutagenesis of an antibody variable domain.

Example 5: Expression of FGF-L Polypeptide in Transgenic Mice

To assess the biological activity of FGF-L polypeptide, a construct encoding an FGF-L polypeptide/Fc fusion protein under the control of a liver specific ApoE promoter is prepared. The delivery of this construct is expected to cause pathological changes that are informative as to the function of FGF-L polypeptide. Similarly, a construct containing the full-length FGF-L polypeptide under the control of the beta actin promoter is prepared. The delivery of this construct is expected to result in ubiquitous expression.

To generate these constructs, PCR is used to amplify template DNA sequences encoding an FGF-L polypeptide using primers that correspond to the 5' and 3' ends of the desired sequence and which incorporate restriction enzyme sites to permit insertion of the amplified product into an expression
5 vector. Following amplification, PCR products are gel purified, digested with the appropriate restriction enzymes, and ligated into an expression vector using standard recombinant DNA techniques. For example, amplified FGF-L polypeptide sequences can be cloned into an expression vector under the control of the human β -actin promoter as described by Graham *et al.*, 1997,
10 *Nature Genetics*, 17:272-74 and Ray *et al.*, 1991, *Genes Dev.* 5:2265-73.

Following ligation, reaction mixtures are used to transform an *E. coli* host strain by electroporation and transformants are selected for drug resistance. Plasmid DNA from selected colonies is isolated and subjected to DNA sequencing to confirm the presence of an appropriate insert and absence
15 of mutation. The FGF-L polypeptide expression vector is purified through two rounds of CsCl density gradient centrifugation, cleaved with a suitable restriction enzyme, and the linearized fragment containing the FGF-L polypeptide transgene is purified by gel electrophoresis. The purified fragment is resuspended in 5 mM Tris, pH 7.4, and 0.2 mM EDTA at a
20 concentration of 2 mg/mL.

Single-cell embryos from BDF1 x BDF1 bred mice are injected as described (PCT Pub. No. WO 97/23614). Embryos are cultured overnight in a CO₂ incubator and 15-20 two-cell embryos are transferred to the oviducts of a pseudopregnant CD1 female mice. Offspring obtained from the implantation
25 of microinjected embryos are screened by PCR amplification of the integrated transgene in genomic DNA samples as follows. Ear pieces are digested in 20 mL ear buffer (20 mM Tris, pH 8.0, 10 mM EDTA, 0.5% SDS, and 500 mg/mL proteinase K) at 55°C overnight. The sample is then diluted with 200 mL of TE, and 2 mL of the ear sample is used in a PCR reaction using
30 appropriate primers.

At 8 weeks of age, transgenic founder animals and control animals are sacrificed for necropsy and pathological analysis. Portions of spleen are

removed and total cellular RNA isolated from the spleens using the Total RNA Extraction Kit (Qiagen) and transgene expression determined by RT-PCR. RNA recovered from spleens is converted to cDNA using the SuperScript™ Preamplification System (Gibco-BRL) as follows. A suitable
5 primer, located in the expression vector sequence and 3' to the FGF-L polypeptide transgene, is used to prime cDNA synthesis from the transgene transcripts. Ten mg of total spleen RNA from transgenic founders and controls is incubated with 1 mM of primer for 10 minutes at 70°C and placed on ice. The reaction is then supplemented with 10 mM Tris-HCl, pH 8.3, 50
10 mM KCl, 2.5 mM MgCl₂, 10 mM of each dNTP, 0.1 mM DTT, and 200 U of SuperScript II reverse transcriptase. Following incubation for 50 minutes at 42°C, the reaction is stopped by heating for 15 minutes at 72°C and digested with 2U of RNase H for 20 minutes at 37°C. Samples are then amplified by PCR using primers specific for FGF-L polypeptide.

15

Example 6: Biological Activity of FGF-L Polypeptide in Transgenic Mice

Prior to euthanasia, transgenic animals are weighed, anesthetized by isofluorane and blood drawn by cardiac puncture. The samples are subjected to hematology and serum chemistry analysis. Radiography is performed after
20 terminal exsanguination. Upon gross dissection, major visceral organs are subject to weight analysis.

Following gross dissection, tissues (*i.e.*, liver, spleen, pancreas, stomach, the entire gastrointestinal tract, kidney, reproductive organs, skin and mammary glands, bone, brain, heart, lung, thymus, trachea, esophagus,
25 thyroid, adrenals, urinary bladder, lymph nodes and skeletal muscle) are removed and fixed in 10% buffered Zn-Formalin for histological examination. After fixation, the tissues are processed into paraffin blocks, and 3 mm sections are obtained. All sections are stained with hematoxylin and eosin, and are then subjected to histological analysis.

30 The spleen, lymph node, and Peyer's patches of both the transgenic and the control mice are subjected to immunohistology analysis with B cell and T cell specific antibodies as follows. The formalin fixed paraffin

embedded sections are deparaffinized and hydrated in deionized water. The sections are quenched with 3% hydrogen peroxide, blocked with Protein Block (Lipshaw, Pittsburgh, PA), and incubated in rat monoclonal anti-mouse B220 and CD3 (Harlan, Indianapolis, IN). Antibody binding is detected by
5 biotinylated rabbit anti-rat immunoglobulins and peroxidase conjugated streptavidin (BioGenex, San Ramon, CA) with DAB as a chromagen (BioTek, Santa Barbara, CA). Sections are counterstained with hematoxylin.

After necropsy, MLN and sections of spleen and thymus from transgenic animals and control littermates are removed. Single cell
10 suspensions are prepared by gently grinding the tissues with the flat end of a syringe against the bottom of a 100 mm nylon cell strainer (Becton Dickinson, Franklin Lakes, NJ). Cells are washed twice, counted, and approximately 1×10^6 cells from each tissue are then incubated for 10 minutes with $0.5 \mu\text{g}$ CD16/32(Fc γ III/II) Fc block in a $20 \mu\text{L}$ volume. Samples are then stained for
15 30 minutes at $2-8^\circ\text{C}$ in a $100 \mu\text{L}$ volume of PBS (lacking Ca^+ and Mg^+), 0.1% bovine serum albumin, and 0.01% sodium azide with $0.5 \mu\text{g}$ antibody of FITC or PE-conjugated monoclonal antibodies against CD90.2 (Thy-1.2), CD45R (B220), CD11b(Mac-1), Gr-1, CD4, or CD8 (PharMingen, San Diego, CA). Following antibody binding, the cells are washed and then analyzed by flow
20 cytometry on a FACScan (Becton Dickinson).

While the present invention has been described in terms of the preferred embodiments, it is understood that variations and modifications will occur to those skilled in the art. Therefore, it is intended that the appended
25 claims cover all such equivalent variations that come within the scope of the invention as claimed.

WHAT IS CLAIMED IS:

1. An isolated nucleic acid molecule comprising a nucleotide sequence selected from the group consisting of:
 - 5 (a) the nucleotide sequence as set forth in SEQ ID NO: 1;
 - (b) the nucleotide sequence of the DNA insert in ATCC Deposit No. PTA-1882;
 - (c) a nucleotide sequence encoding the polypeptide as set forth in SEQ ID NO: 2;
 - 10 (d) a nucleotide sequence which hybridizes under moderately or highly stringent conditions to the complement of any of (a) - (c); and
 - (e) a nucleotide sequence complementary to any of (a) - (c).
2. An isolated nucleic acid molecule comprising a nucleotide
15 sequence selected from the group consisting of:
 - (a) a nucleotide sequence encoding a polypeptide which is at least about 70 percent identical to the polypeptide as set forth in SEQ ID NO: 2, wherein the encoded polypeptide has an activity of the polypeptide set forth in SEQ ID NO: 2;
 - 20 (b) a nucleotide sequence encoding an allelic variant or splice variant of the nucleotide sequence as set forth in SEQ ID NO: 1, the nucleotide sequence of the DNA insert in ATCC Deposit No. PTA-1882, or (a);
 - (c) a region of the nucleotide sequence of SEQ ID NO: 1, the DNA
25 insert in ATCC Deposit No. PTA-1882, (a), or (b) encoding a polypeptide fragment of at least about 25 amino acid residues, wherein the polypeptide fragment has an activity of the encoded polypeptide as set forth in SEQ ID NO: 2, or is antigenic;
 - (d) a region of the nucleotide sequence of SEQ ID NO: 1, the DNA
30 insert in ATCC Deposit No. PTA-1882, or any of (a) - (c) comprising a fragment of at least about 16 nucleotides;

(e) a nucleotide sequence which hybridizes under moderately or highly stringent conditions to the complement of any of (a) - (d); and

(f) a nucleotide sequence complementary to any of (a) - (d).

5 3. An isolated nucleic acid molecule comprising a nucleotide sequence selected from the group consisting of:

(a) a nucleotide sequence encoding a polypeptide as set forth in SEQ ID NO: 2 with at least one conservative amino acid substitution, wherein the encoded polypeptide has an activity of the polypeptide set forth in SEQ ID

10 NO: 2;

(b) a nucleotide sequence encoding a polypeptide as set forth in SEQ ID NO: 2 with at least one amino acid insertion, wherein the encoded polypeptide has an activity of the polypeptide set forth in SEQ ID NO: 2;

(c) a nucleotide sequence encoding a polypeptide as set forth in
15 SEQ ID NO: 2 with at least one amino acid deletion, wherein the encoded polypeptide has an activity of the polypeptide set forth in SEQ ID NO: 2;

(d) a nucleotide sequence encoding a polypeptide as set forth in SEQ ID NO: 2 which has a C- and/or N- terminal truncation, wherein the encoded polypeptide has an activity of the polypeptide set forth in SEQ ID
20 NO: 2;

(e) a nucleotide sequence encoding a polypeptide as set forth in SEQ ID NO: 2 with at least one modification selected from the group consisting of amino acid substitutions, amino acid insertions, amino acid deletions, C-terminal truncation, and N-terminal truncation, wherein the
25 encoded polypeptide has an activity of the polypeptide set forth in SEQ ID NO: 2;

(f) a nucleotide sequence of any of (a) - (e) comprising a fragment of at least about 16 nucleotides;

(g) a nucleotide sequence which hybridizes under moderately or
30 highly stringent conditions to the complement of any of (a) - (f); and

(h) a nucleotide sequence complementary to any of (a) - (e).

4. A vector comprising the nucleic acid molecule of any of Claims 1, 2, or 3.
5. A host cell comprising the vector of Claim 4.
6. The host cell of Claim 5 that is a eukaryotic cell.
7. The host cell of Claim 5 that is a prokaryotic cell.
8. A process of producing an FGF-L polypeptide comprising culturing the host cell of Claim 5 under suitable conditions to express the polypeptide, and optionally isolating the polypeptide from the culture.
9. A polypeptide produced by the process of Claim 8.
10. The process of Claim 8, wherein the nucleic acid molecule comprises promoter DNA other than the promoter DNA for the native FGF-L polypeptide operatively linked to the DNA encoding the FGF-L polypeptide.
11. The isolated nucleic acid molecule according to Claim 2, wherein the percent identity is determined using a computer program selected from the group consisting of GAP, BLASTN, FASTA, BLASTA, BLASTX, BestFit, and the Smith-Waterman algorithm.
12. A process for determining whether a compound inhibits FGF-L polypeptide activity or FGF-L polypeptide production comprising exposing a cell according to any of Claims 5, 6, or 7 to the compound and measuring FGF-L polypeptide activity or FGF-L polypeptide production in said cell.
13. An isolated polypeptide comprising the amino acid sequence selected from the group consisting of:
- (a) the amino acid sequence as set forth in SEQ ID NO: 2; and

(b) the amino acid sequence encoded by the DNA insert in ATCC Deposit No. PTA-1882.

14. An isolated polypeptide comprising the amino acid sequence
5 selected from the group consisting of:
- (a) an amino acid sequence for an ortholog of SEQ ID NO: 2;
 - (b) an amino acid sequence which is at least about 70 percent identical to the amino acid sequence of SEQ ID NO: 2, wherein the polypeptide has an activity of the polypeptide set forth in SEQ ID NO: 2;
 - 10 (c) a fragment of the amino acid sequence set forth in SEQ ID NO: 2 comprising at least about 25 amino acid residues, wherein the fragment has an activity of the polypeptide set forth in SEQ ID NO: 2, or is antigenic; and
 - (d) an amino acid sequence for an allelic variant or splice variant of the amino acid sequence as set forth in SEQ ID NO: 2, the amino acid
15 sequence encoded by the DNA insert in ATCC Deposit No. PTA-1882, (a), or (b).

15. An isolated polypeptide comprising the amino acid sequence selected from the group consisting of:
- 20 (a) the amino acid sequence as set forth in SEQ ID NO: 2 with at least one conservative amino acid substitution, wherein the polypeptide has an activity of the polypeptide set forth in SEQ ID NO: 2;
 - (b) the amino acid sequence as set forth in SEQ ID NO: 2 with at least one amino acid insertion, wherein the polypeptide has an activity of the
25 polypeptide set forth in SEQ ID NO: 2;
 - (c) the amino acid sequence as set forth in SEQ ID NO: 2 with at least one amino acid deletion, wherein the polypeptide has an activity of the polypeptide set forth in SEQ ID NO: 2;
 - (d) the amino acid sequence as set forth in SEQ ID NO: 2 which
30 has a C- and/or N- terminal truncation, wherein the polypeptide has an activity of the polypeptide set forth in SEQ ID NO: 2; and

(e) the amino acid sequence as set forth in SEQ ID NO: 2 with at least one modification selected from the group consisting of amino acid substitutions, amino acid insertions, amino acid deletions, C-terminal truncation, and N-terminal truncation, wherein the polypeptide has an activity of the polypeptide set forth in SEQ ID NO: 2.

16. An isolated polypeptide encoded by the nucleic acid molecule of any of Claims 1, 2, or 3, wherein the polypeptide has an activity of the polypeptide set forth in SEQ ID NO: 2.

17. The isolated polypeptide according to Claim 14, wherein the percent identity is determined using a computer program selected from the group consisting of GAP, BLASTP, FASTA, BLASTA, BLASTX, BestFit, and the Smith-Waterman algorithm.

18. A selective binding agent or fragment thereof which specifically binds the polypeptide of any of Claims 13, 14, or 15.

19. The selective binding agent or fragment thereof of Claim 18 that specifically binds the polypeptide comprising the amino acid sequence as set forth in SEQ ID NO: 2, or a fragment thereof.

20. The selective binding agent of Claim 18 that is an antibody or fragment thereof.

21. The selective binding agent of Claim 18 that is a humanized antibody.

22. The selective binding agent of Claim 18 that is a human antibody or fragment thereof.

23. The selective binding agent of Claim 18 that is a polyclonal antibody or fragment thereof.

24. The selective binding agent Claim 18 that is a monoclonal
5 antibody or fragment thereof.

25. The selective binding agent of Claim 18 that is a chimeric antibody or fragment thereof.

10 26. The selective binding agent of Claim 18 that is a CDR-grafted antibody or fragment thereof.

27. The selective binding agent of Claim 18 that is an antiidiotypic antibody or fragment thereof.

15

28. The selective binding agent of Claim 18 that is a variable region fragment.

29. The variable region fragment of Claim 28 that is a Fab or a
20 Fab' fragment.

30. A selective binding agent or fragment thereof comprising at least one complementarity determining region with specificity for a polypeptide having the amino acid sequence of SEQ ID NO: 2.

25

31. The selective binding agent of Claim 18 that is bound to a detectable label.

32. The selective binding agent of Claim 18 that antagonizes FGF-
30 L polypeptide biological activity.

33. A method for treating, preventing, or ameliorating an FGF-L polypeptide-related disease, condition, or disorder comprising administering to a patient an effective amount of a selective binding agent according to Claim 18.

5

34. A selective binding agent produced by immunizing an animal with a polypeptide comprising an amino acid sequence of SEQ ID NO: 2.

35. A hybridoma which produces a selective binding agent which
10 is capable of binding a polypeptide according to any of Claims 1, 2, or 3.

36. A method of detecting or quantitating the amount of FGF-L polypeptide using the selective binding agent or fragment of Claim 18.

15 37. A composition comprising the polypeptide of any of Claims 13, 14, or 15, and a pharmaceutically acceptable formulation agent.

38. The composition of Claim 37, wherein the pharmaceutically acceptable formulation agent is a carrier, adjuvant, solubilizer, stabilizer, or
20 anti-oxidant.

39. A polypeptide comprising a derivative of the polypeptide of any of Claims 13, 14, or 15.

25 40. The polypeptide of Claim 39 that is covalently modified with a water-soluble polymer.

41. The polypeptide of Claim 40, wherein the water-soluble polymer is selected from the group consisting of polyethylene glycol, monomethoxy-polyethylene glycol, dextran, cellulose, poly-(N-vinyl pyrrolidone) polyethylene glycol, propylene glycol homopolymers,
30

polypropylene oxide/ethylene oxide co-polymers, polyoxyethylated polyols, and polyvinyl alcohol.

42. A composition comprising a nucleic acid molecule of any of
5 Claims 1, 2, or 3 and a pharmaceutically acceptable formulation agent.

43. The composition of Claim 42, wherein said nucleic acid molecule is contained in a viral vector.

10 44. A viral vector comprising a nucleic acid molecule of any of Claims 1, 2, or 3.

45. A fusion polypeptide comprising the polypeptide of any of Claims 13, 14, or 15 fused to a heterologous amino acid sequence.

15

46. The fusion polypeptide of Claim 45, wherein the heterologous amino acid sequence is an IgG constant domain or fragment thereof.

47. A method for treating, preventing, or ameliorating a medical
20 condition comprising administering to a patient the polypeptide of any of Claims 13, 14, or 15, or the polypeptide encoded by the nucleic acid of any of Claims 1, 2, or 3.

48. A method of diagnosing a pathological condition or a
25 susceptibility to a pathological condition in a subject comprising:

(a) determining the presence or amount of expression of the polypeptide of any of Claims 13, 14, or 15, or the polypeptide encoded by the nucleic acid molecule of any of Claims 1, 2, or 3 in a sample; and

(b) diagnosing a pathological condition or a susceptibility to a
30 pathological condition based on the presence or amount of expression of the polypeptide.

49. A device, comprising:
(a) a membrane suitable for implantation; and
(b) cells encapsulated within said membrane, wherein said cells secrete a protein of any of Claims 13, 14, or 15; and

5 said membrane is permeable to said protein and impermeable to materials detrimental to said cells.

50. A method of identifying a compound which binds to an FGF-L polypeptide comprising:

10 (a) contacting the polypeptide of any of Claims 13, 14, or 15 with a compound; and

 (b) determining the extent of binding of the FGF-L polypeptide to the compound.

15 51. The method of Claim 50, further comprising determining the activity of the polypeptide when bound to the compound.

52. A method of modulating levels of a polypeptide in an animal comprising administering to the animal the nucleic acid molecule of any of
20 Claims 1, 2, or 3.

53. A transgenic non-human mammal comprising the nucleic acid molecule of any of Claims 1, 2, or 3.

25 54. A process for determining whether a compound inhibits FGF-L polypeptide activity or FGF-L polypeptide production comprising exposing a transgenic mammal according to Claim 53 to the compound, and measuring FGF-L polypeptide activity or FGF-L polypeptide production in said mammal.

FIG. 1A

```

tacacggccg cagctgaaca gcatcacgc tgtcccaagg acaaccccaa agaggggcct 60
cgactgcacc tcctcgaagt tgctggctgg ctttggaag tgcaggaatg gtgttttgtg 120
agggcatgga tggagaagtg ccaagggccc ctgtttggtc acttccgaag agcaaaaacg 180
tgttgagagg agaccggttt aagatttcaa acagaacctc cccagcgcgc atgaaaggac 240
ttgattagca tatgtcaaga ggacccgctt atatactcg tgtgtatgta cacaggactc 300
tgatctgac agtttgcgga attggagccc cagccaacag ccctagtcct agtattggca 360
gcggcagcta tagatatttc tgcagagcca gcagccggct cccacctacc caaggagaga 420
agatcgctcc aagacagtga gagcttcctt gccatttcag tgcaaagtcc ctccggagcg 480
acctcagagg agtaaccggg ccttaacttt ttgcgctcgt tttgctataa tttttctcta 540
tccacctcca tcccaccccc acaacactct ttactggggg ggtcttttgt gttccggatc 600
tccccctcc atg gct ccc tta gcc gaa gtc ggg ggc ttt ctg ggc ggc ctg 651
      Met Ala Pro Leu Ala Glu Val Gly Gly Phe Leu Gly Gly Leu
        1             5             10

gag ggc ttg ggc cag cag gtg ggt tcg cat ttc ctg ttg cct cct gcc 699
Glu Gly Leu Gly Gln Gln Val Gly Ser His Phe Leu Leu Pro Pro Ala
  15             20             25             30

ggg gag cgg ccg ccg ctg ctg ggc gag cgc agg agc gcg gcg gag cgg 747
Gly Glu Arg Pro Pro Leu Leu Gly Glu Arg Arg Ser Ala Ala Glu Arg
      35             40             45

agc gcc cgc ggc ggg ccg ggg gct gcg cag ctg gcg cac ctg cac ggc 795
Ser Ala Arg Gly Gly Pro Gly Ala Ala Gln Leu Ala His Leu His Gly
      50             55             60

atc ctg cgc cgc cgg cag ctc tat tgc cgc acc ggc ttc cac ctg cag 843
Ile Leu Arg Arg Arg Gln Leu Tyr Cys Arg Thr Gly Phe His Leu Gln
      65             70             75

atc ctg ccc gac ggc agc gtg cag ggc acc cgg cag gac cac agc ctc 891
Ile Leu Pro Asp Gly Ser Val Gln Gly Thr Arg Gln Asp His Ser Leu
      80             85             90

ttc ggt atc ttg gaa ttc atc agt gtg gca gtg gga ctg gtc agt att 939
Phe Gly Ile Leu Glu Phe Ile Ser Val Ala Val Gly Leu Val Ser Ile
      95             100             105             110

aga ggt gtg gac agt ggt ctc tat ctt gga atg aat gac aaa gga gaa 987
Arg Gly Val Asp Ser Gly Leu Tyr Leu Gly Met Asn Asp Lys Gly Glu
      115             120             125

```

FIG. 1B

ctc tat gga tca gag aaa ctt act tcc gaa tgc atc ttt agg gag cag	1035
Leu Tyr Gly Ser Glu Lys Leu Thr Ser Glu Cys Ile Phe Arg Glu Gln	
130 135 140	
ttt gaa gag aac tgg tat aac acc tat tca tct aac ata tat aaa cat	1083
Phe Glu Glu Asn Trp Tyr Asn Thr Tyr Ser Ser Asn Ile Tyr Lys His	
145 150 155	
gga gac act ggc cgc agg tat ttt gtg gca ctt aac aaa gac gga act	1131
Gly Asp Thr Gly Arg Arg Tyr Phe Val Ala Leu Asn Lys Asp Gly Thr	
160 165 170	
cca aga gat ggc gcc agg tcc aag agg cat cag aaa ttt aca cat ttc	1179
Pro Arg Asp Gly Ala Arg Ser Lys Arg His Gln Lys Phe Thr His Phe	
175 180 185 190	
tta cct aga cca gtg gat cca gaa aga gtt cca gaa ttg tac aag gac	1227
Leu Pro Arg Pro Val Asp Pro Glu Arg Val Pro Glu Leu Tyr Lys Asp	
195 200 205	
cta ctg atg tac act tga agtgcgatag tgacattatg gaagagtcaa	1275
Leu Leu Met Tyr Thr	
210	
accacaacca ttctttcttg tcatagttcc catcataaaa taatgaccca agcag	1330

[illegible]

FIG. 3

```
1 MAPLAEVGGFLGGLEGLGQQVGSFLLPPAGERPPLLGERRSAAERSA.R 49
  |||| ||| : | : . |. .|| : || : .|
1 MAPLGEVGSYFGVQDAV..PFGNVPVLPV..DSPVLLSDHLGQSEAGGLP 46

50 GGPGAAQLAHLHGILRRRQLYCRTGFHLQILPDGSVQGTRQDHSLEFGILE 99
  || | || ||||| ||||| : | |. : |||. || ||||
47 RGPVAVTDLHLKGILRRRQLYCRTGFHLEIFPNGTIQGTRKDHRSRFGILE 96

100 FISVAVGLVSIRGVDSGLYLGMDKGEYLGSEKLTSECIFREQFEENWYN 149
  ||| : ||||| ||||| ||||| : ||||| ||||| ||| : ||||| |||||
97 FISIAVGLVSIRGVDSGLYLGMDKGEYLGSEKLTQECVFEENWYN 146

150 TYSSNIYKHGDTGRRYFVALNKDGTPRDGARSKRHQKFTHFLPRPVDPER 199
  |||| : ||| ||||| : ||||| ||||| : | |. ||||| ||||| ||||| : :
147 TYSSNLYKHVDTGRRYYVALNKDGTREGTRTKRHQKFTHFLPRPVDPAK 196

200 VPELYKDLLMYT 211
  ||||| : | .
197 VPELYKDILSQS 208
```

FIG. 4

hu FGF-9	MAPLGEVGNV	FGVQDAV..P	FGNVPLPV.	.DSPVLLSDH	LQSEAGGLP	RGPAVTDLDH
hu FGF-16	MA...EVGGV	FASLDWDLHG	FSSSLGNVPL	ADSPGFLNER	LGQIE.GKLQ	RGSP.TDFAH
hu FGF-L	MAPLAEVGGF	LGLEGLGQQ	VGSHFLPPA	GERPPLLGER	RSAAERSA.R	GGPGAAQLAH
mu FGF-9	MAPLGEVGSY	FGVQDAV..P	FGNVPLPV.	.DSPVLLNDH	LQSEAGGLP	RGPAVTDLDH
ra FGF-16	MA...EVGGV	FASLDWDLQG	FSSSLGNVPL	ADSPGFLNER	LGQIE.GKLQ	RGSP.TDFAH
hu FGF-9	LKGILRRRQL	YCRTGFHLEI	FPNGTIQCTR	KHSRFGILE	FISIAVGLVS	IRGVDSGLYL
hu FGF-16	LKGILRRRQL	YCRTGFHLEI	FPNGTVHGTR	HDHSRFGILE	FISLAVGLIS	IRGVDSGLYL
hu FGF-L	LHGILRRRQL	YCRTGFHLQI	LPDGSVQCTR	QDHSRFGILE	FISVAVGLVS	IRGVDSGLYL
mu FGF-9	LKGILRRRQL	YCRTGFHLEI	FPNGTIQCTR	KHSRFGILE	FISIAVGLVS	IRGVDSGLYL
ra FGF-16	LKGILRRRQL	YCRTGFHLEI	FPNGTVHGTR	HDHSRFGILE	FISLAVGLIS	IRGVDSGLYL
hu FGF-9	GMNEKGELYG	SEKLTQECVF	REQFEENWYN	TYSSNLYKHV	DTGRRYYVAL	NKDGTPREGT
hu FGF-16	GMNERGELYG	SKKLTRECVF	REQFEENWYN	TYASTLYKHS	DSERQYYVAL	NKDGSPREGY
hu FGF-L	GMNDKGELYG	SEKLTSECIF	REQFEENWYN	TYSSNIYKHG	DTGRRYFVAL	NKDGTPRDGA
mu FGF-9	GMNEKGELYG	SEKLTQECVF	REQFEENWYN	TYSSNLYKHV	DTGRRYYVAL	NKDGTPREGT
ra FGF-16	GMNERGELFG	SKKLTRECVF	REQFEENWYN	TYASTLYKHS	DSERQYYVAL	NKDGSPREGY
hu FGF-9	RTKRHHQKPTH	FLPRPVDPK	VPELYKDILS	QS		
hu FGF-16	RTKRHHQKPTH	FLPRPVDPSK	LPSMSRDLFH	YR		
hu FGF-L	RSKRHHQKPTH	FLPRPVDPER	VPELYKDILM	YT		
mu FGF-9	RTKRHHQKPTH	FLPRPVDPK	VPELYKDILS	QS		
ra FGF-16	RTKRHHQKPTH	FLPRPVDPSK	LPSMSRDLFR	YR		

SEQUENCE LISTING

<110> Bass, Michael B.
Jing, Shuqian

<120> Fibroblast Growth Factor-Like Molecules and Uses
Thereof

<130> 01-006-A1

<140>

<141>

<150> 60/188,786

<151> 2000-03-13

<160> 11

<170> PatentIn Ver. 2.0

<210> 1

<211> 1330

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (610)..(1245)

<400> 1

```

tacacggccg cagctgaaca gcatcaccgc tgtcccaagg acaaccccaa agaggggcct 60
cgactgcacc tctcgaagt tgctggctgg ctttggcaag tgcaggaatg gtgttttgtg 120
agggcatgga tggagaagt ccaagggccc ctgtttggtc acttccgaag agcaaaaacg 180
tgttgagagg agaccggttt aagatttcaa acagaacctc cccagcgcgc atgaaaggac 240
ttgattagca tatgtcaaga ggaccgcgtt atatactcgg tgtgtatgta cacaggactc 300
tgatctgata agtttgcgga attggagccc cagccaacag ccctagtcct agtattggca 360
gcggcagcta tagatatctc tgcagagcca gcagccggct cccacctacc caaggagaga 420
agatcgctcc aagacagtga gagcttcctt gccatttcag tgcaaagtcc ctccggagcg 480
acctcagagg agtaaccggg ccttaacttt ttgcgctcgt ttgtctataa tttttctcta 540
tccacctcca tcccacccc acaacactct ttactggggg ggtcttttgt gttccggatc 600
tccccctcc atg gct ccc tta gcc gaa gtc ggg ggc ttt ctg ggc ggc ctg 651
      Met Ala Pro Leu Ala Glu Val Gly Gly Phe Leu Gly Gly Leu
        1              5              10

gag ggc ttg ggc cag cag gtg ggt tcg cat ttc ctg ttg cct cct gcc 699
Glu Gly Leu Gly Gln Gln Val Gly Ser His Phe Leu Leu Pro Pro Ala
  15              20              25              30

```

```

ggg gag cgg ccg ccg ctg ctg ggc gag cgc agg agc gcg gcg gag cgg 747
Gly Glu Arg Pro Pro Leu Leu Gly Glu Arg Arg Ser Ala Ala Glu Arg
          35          40          45

agc gcc cgc ggc ggg ccg ggg gct gcg cag ctg gcg cac ctg cac ggc 795
Ser Ala Arg Gly Gly Pro Gly Ala Ala Gln Leu Ala His Leu His Gly
          50          55          60

atc ctg cgc cgc cgg cag ctc tat tgc cgc acc ggc ttc cac ctg cag 843
Ile Leu Arg Arg Arg Gln Leu Tyr Cys Arg Thr Gly Phe His Leu Gln
          65          70          75

atc ctg ccc gac ggc agc gtg cag ggc acc cgg cag gac cac agc ctc 891
Ile Leu Pro Asp Gly Ser Val Gln Gly Thr Arg Gln Asp His Ser Leu
          80          85          90

ttc ggt atc ttg gaa ttc atc agt gtg gca gtg gga ctg gtc agt att 939
Phe Gly Ile Leu Glu Phe Ile Ser Val Ala Val Gly Leu Val Ser Ile
          95          100          105          110

aga ggt gtg gac agt ggt ctc tat ctt gga atg aat gac aaa gga gaa 987
Arg Gly Val Asp Ser Gly Leu Tyr Leu Gly Met Asn Asp Lys Gly Glu
          115          120          125

ctc tat gga tca gag aaa ctt act tcc gaa tgc atc ttt agg gag cag 1035
Leu Tyr Gly Ser Glu Lys Leu Thr Ser Glu Cys Ile Phe Arg Glu Gln
          130          135          140

ttt gaa gag aac tgg tat aac acc tat tca tct aac ata tat aaa cat 1083
Phe Glu Glu Asn Trp Tyr Asn Thr Tyr Ser Ser Asn Ile Tyr Lys His
          145          150          155

gga gac act ggc cgc agg tat ttt gtg gca ctt aac aaa gac gga act 1131
Gly Asp Thr Gly Arg Arg Tyr Phe Val Ala Leu Asn Lys Asp Gly Thr
          160          165          170

cca aga gat ggc gcc agg tcc aag agg cat cag aaa ttt aca cat ttc 1179
Pro Arg Asp Gly Ala Arg Ser Lys Arg His Gln Lys Phe Thr His Phe
          175          180          185          190

tta cct aga cca gtg gat cca gaa aga gtt cca gaa ttg tac aag gac 1227
Leu Pro Arg Pro Val Asp Pro Glu Arg Val Pro Glu Leu Tyr Lys Asp
          195          200          205

cta ctg atg tac act tga agtgcgatag tgacattatg gaagagtcaa 1275
Leu Leu Met Tyr Thr
          210

accacaacca ttctttcttg tcatagttcc catcataaaa taatgaccca agcag 1330

<210> 2
<211> 211
<212> PRT
<213> Homo sapiens

<400> 2
Met Ala Pro Leu Ala Glu Val Gly Gly Phe Leu Gly Gly Leu Glu Gly

```

1					5					10					15				
Leu	Gly	Gln	Gln	Val	Gly	Ser	His	Phe	Leu	Leu	Pro	Pro	Ala	Gly	Glu				
			20							25				30					
Arg	Pro	Pro	Leu	Leu	Gly	Glu	Arg	Arg	Ser	Ala	Ala	Glu	Arg	Ser	Ala				
			35							40				45					
Arg	Gly	Gly	Pro	Gly	Ala	Ala	Gln	Leu	Ala	His	Leu	His	Gly	Ile	Leu				
			50							55				60					
Arg	Arg	Arg	Gln	Leu	Tyr	Cys	Arg	Thr	Gly	Phe	His	Leu	Gln	Ile	Leu				
			65							70				75					
Pro	Asp	Gly	Ser	Val	Gln	Gly	Thr	Arg	Gln	Asp	His	Ser	Leu	Phe	Gly				
			85							90				95					
Ile	Leu	Glu	Phe	Ile	Ser	Val	Ala	Val	Gly	Leu	Val	Ser	Ile	Arg	Gly				
			100							105				110					
Val	Asp	Ser	Gly	Leu	Tyr	Leu	Gly	Met	Asn	Asp	Lys	Gly	Glu	Leu	Tyr				
			115							120				125					
Gly	Ser	Glu	Lys	Leu	Thr	Ser	Glu	Cys	Ile	Phe	Arg	Glu	Gln	Phe	Glu				
			130							135				140					
Glu	Asn	Trp	Tyr	Asn	Thr	Tyr	Ser	Ser	Asn	Ile	Tyr	Lys	His	Gly	Asp				
			145							150				155					
Thr	Gly	Arg	Arg	Tyr	Phe	Val	Ala	Leu	Asn	Lys	Asp	Gly	Thr	Pro	Arg				
			165							170				175					
Asp	Gly	Ala	Arg	Ser	Lys	Arg	His	Gln	Lys	Phe	Thr	His	Phe	Leu	Pro				
			180							185				190					
Arg	Pro	Val	Asp	Pro	Glu	Arg	Val	Pro	Glu	Leu	Tyr	Lys	Asp	Leu	Leu				
			195							200				205					
Met	Tyr	Thr																	
			210																

```
<210> 3
<211> 208
<212> PRT
<213> Homo sapiens
```

```

<400> 3
Met Ala Pro Leu Gly Glu Val Gly Asn Tyr Phe Gly Val Gln Asp Ala
  1             5             10             15
Val Pro Phe Gly Asn Val Pro Val Leu Pro Val Asp Ser Pro Val Leu
          20             25             30
Leu Ser Asp His Leu Gly Gln Ser Glu Ala Gly Gly Leu Pro Arg Gly
      35             40             45
Pro Ala Val Thr Asp Leu Asp His Leu Lys Gly Ile Leu Arg Arg Arg

```

50 55 60
 Gln Leu Tyr Cys Arg Thr Gly Phe His Leu Glu Ile Phe Pro Asn Gly
 65 70 75 80
 Thr Ile Gln Gly Thr Arg Lys Asp His Ser Arg Phe Gly Ile Leu Glu
 85 90 95
 Phe Ile Ser Ile Ala Val Gly Leu Val Ser Ile Arg Gly Val Asp Ser
 100 105 110
 Gly Leu Tyr Leu Gly Met Asn Glu Lys Gly Glu Leu Tyr Gly Ser Glu
 115 120 125
 Lys Leu Thr Gln Glu Cys Val Phe Arg Glu Gln Phe Glu Glu Asn Trp
 130 135 140
 Tyr Asn Thr Tyr Ser Ser Asn Leu Tyr Lys His Val Asp Thr Gly Arg
 145 150 155 160
 Arg Tyr Tyr Val Ala Leu Asn Lys Asp Gly Thr Pro Arg Glu Gly Thr
 165 170 175
 Arg Thr Lys Arg His Gln Lys Phe Thr His Phe Leu Pro Arg Pro Val
 180 185 190
 Asp Pro Asp Lys Val Pro Glu Leu Tyr Lys Asp Ile Leu Ser Gln Ser
 195 200 205

<210> 4
 <211> 208
 <212> PRT
 <213> Rattus norvegicus

<400> 4
 Met Ala Pro Leu Gly Glu Val Gly Ser Tyr Phe Gly Val Gln Asp Ala
 1 5 10 15
 Val Pro Phe Gly Asn Val Pro Val Leu Pro Val Asp Ser Pro Val Leu
 20 25 30
 Leu Ser Asp His Leu Gly Gln Ser Glu Ala Gly Gly Leu Pro Arg Gly
 35 40 45
 Pro Ala Val Thr Asp Leu Asp His Leu Lys Gly Ile Leu Arg Arg Arg
 50 55 60
 Gln Leu Tyr Cys Arg Thr Gly Phe His Leu Glu Ile Phe Pro Asn Gly
 65 70 75 80
 Thr Ile Gln Gly Thr Arg Lys Asp His Ser Arg Phe Gly Ile Leu Glu
 85 90 95
 Phe Ile Ser Ile Ala Val Gly Leu Val Ser Ile Arg Gly Val Asp Ser

```

          100              105              110
Gly Leu Tyr Leu Gly Met Asn Glu Lys Gly Glu Leu Tyr Gly Ser Glu
      115              120              125
Lys Leu Thr Gln Glu Cys Val Phe Arg Glu Gln Phe Glu Glu Asn Trp
      130              135              140
Tyr Asn Thr Tyr Ser Ser Asn Leu Tyr Lys His Val Asp Thr Gly Arg
      145              150              155              160
Arg Tyr Tyr Val Ala Leu Asn Lys Asp Gly Thr Pro Arg Glu Gly Thr
      165              170              175
Arg Thr Lys Arg His Gln Lys Phe Thr His Phe Leu Pro Arg Pro Val
      180              185              190
Asp Pro Asp Lys Val Pro Glu Leu Tyr Lys Asp Ile Leu Ser Gln Ser
      195              200              205

```

```

<210> 5
<211> 207
<212> PRT
<213> Homo sapiens

```

```

<400> 5
Met Ala Glu Val Gly Gly Val Phe Ala Ser Leu Asp Trp Asp Leu His
  1              5              10              15
Gly Phe Ser Ser Ser Leu Gly Asn Val Pro Leu Ala Asp Ser Pro Gly
      20              25              30
Phe Leu Asn Glu Arg Leu Gly Gln Ile Glu Gly Lys Leu Gln Arg Gly
      35              40              45
Ser Pro Thr Asp Phe Ala His Leu Lys Gly Ile Leu Arg Arg Arg Gln
      50              55              60
Leu Tyr Cys Arg Thr Gly Phe His Leu Glu Ile Phe Pro Asn Gly Thr
      65              70              75              80
Val His Gly Thr Arg His Asp His Ser Arg Phe Gly Ile Leu Glu Phe
      85              90              95
Ile Ser Leu Ala Val Gly Leu Ile Ser Ile Arg Gly Val Asp Ser Gly
      100              105              110
Leu Tyr Leu Gly Met Asn Glu Arg Gly Glu Leu Tyr Gly Ser Lys Lys
      115              120              125
Leu Thr Arg Glu Cys Val Phe Arg Glu Gln Phe Glu Glu Asn Trp Tyr
      130              135              140
Asn Thr Tyr Ala Ser Thr Leu Tyr Lys His Ser Asp Ser Glu Arg Gln

```

145 150 155 160
 Tyr Tyr Val Ala Leu Asn Lys Asp Gly Ser Pro Arg Glu Gly Tyr Arg
 165 170 175
 Thr Lys Arg His Gln Lys Phe Thr His Phe Leu Pro Arg Pro Val Asp
 180 185 190
 Pro Ser Lys Leu Pro Ser Met Ser Arg Asp Leu Phe His Tyr Arg
 195 200 205

<210> 6
 <211> 208
 <212> PRT
 <213> Mus musculus

<400> 6
 Met Ala Pro Leu Gly Glu Val Gly Ser Tyr Phe Gly Val Gln Asp Ala
 1 5 10 15
 Val Pro Phe Gly Asn Val Pro Val Leu Pro Val Asp Ser Pro Val Leu
 20 25 30
 Leu Asn Asp His Leu Gly Gln Ser Glu Ala Gly Gly Leu Pro Arg Gly
 35 40 45
 Pro Ala Val Thr Asp Leu Asp His Leu Lys Gly Ile Leu Arg Arg Arg
 50 55 60
 Gln Leu Tyr Cys Arg Thr Gly Phe His Leu Glu Ile Phe Pro Asn Gly
 65 70 75 80
 Thr Ile Gln Gly Thr Arg Lys Asp His Ser Arg Phe Gly Ile Leu Glu
 85 90 95
 Phe Ile Ser Ile Ala Val Gly Leu Val Ser Ile Arg Gly Val Asp Ser
 100 105 110
 Gly Leu Tyr Leu Gly Met Asn Glu Lys Gly Glu Leu Tyr Gly Ser Glu
 115 120 125
 Lys Leu Thr Gln Glu Cys Val Phe Arg Glu Gln Phe Glu Glu Asn Trp
 130 135 140
 Tyr Asn Thr Tyr Ser Ser Asn Leu Tyr Lys His Val Asp Thr Gly Arg
 145 150 155 160
 Arg Tyr Tyr Val Ala Leu Asn Lys Asp Gly Thr Pro Arg Glu Gly Thr
 165 170 175
 Arg Thr Lys Arg His Gln Lys Phe Thr His Phe Leu Pro Arg Pro Val
 180 185 190
 Asp Pro Asp Lys Val Pro Glu Leu Tyr Lys Asp Ile Leu Ser Gln Ser
 195 200 205

<210> 7

<211> 207

<212> PRT

<213> Mus musculus

<400> 7

```

Met Ala Glu Val Gly Gly Val Phe Ala Ser Leu Asp Trp Asp Leu Gln
  1             5             10             15

Gly Phe Ser Ser Ser Leu Gly Asn Val Pro Leu Ala Asp Ser Pro Gly
          20             25             30

Phe Leu Asn Glu Arg Leu Gly Gln Ile Glu Gly Lys Leu Gln Arg Gly
          35             40             45

Ser Pro Thr Asp Phe Ala His Leu Lys Gly Ile Leu Arg Arg Arg Gln
          50             55             60

Leu Tyr Cys Arg Thr Gly Phe His Leu Glu Ile Phe Pro Asn Gly Thr
          65             70             75             80

Val His Gly Thr Arg His Asp His Ser Arg Phe Gly Ile Leu Glu Phe
          85             90             95

Ile Ser Leu Ala Val Gly Leu Ile Ser Ile Arg Gly Val Asp Ser Gly
          100            105            110

Leu Tyr Leu Gly Met Asn Glu Arg Gly Glu Leu Phe Gly Ser Lys Lys
          115            120            125

Leu Thr Arg Glu Cys Val Phe Arg Glu Gln Phe Glu Glu Asn Trp Tyr
          130            135            140

Asn Thr Tyr Ala Ser Thr Leu Tyr Lys His Ser Asp Ser Glu Arg Gln
          145            150            155            160

Tyr Tyr Val Ala Leu Asn Lys Asp Gly Ser Pro Arg Glu Gly Tyr Arg
          165            170            175

Thr Lys Arg His Gln Lys Phe Thr His Phe Leu Pro Arg Pro Val Asp
          180            185            190

Pro Ser Lys Leu Pro Ser Met Ser Arg Asp Leu Phe Arg Tyr Arg
          195            200            205

```

<210> 8

<211> 11

<212> PRT

<213> Human immunodeficiency virus type 1

<400> 8

```

Tyr Gly Arg Lys Lys Arg Arg Gln Arg Arg Arg
  1             5             10

```

<210> 9
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: internalizing
domain derived from HIV tat protein

<400> 9
Gly Gly Gly Gly Tyr Gly Arg Lys Lys Arg Arg Gln Arg Arg Arg
1 5 10 15

<210> 10
<211> 23
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:
oligonucleotide; PCR primer 2440-39

<400> 10
ccatggctcc cttagccgaa gtc 23

<210> 11
<211> 26
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:
oligonucleotide; PCR primer 2432-77

<400> 11
tcactatcgc acttcaagtg tacatc 26

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
20 September 2001 (20.09.2001)

PCT

(10) International Publication Number
WO 01/68854 A3

(51) International Patent Classification⁷: C12N 15/12,
C07K 14/50, C12N 1/21, 5/10, C07K 16/22, A61K
39/395, G01N 33/68, A61K 38/18, 31/70, C12N 15/62,
A01K 67/027, C12N 5/22

(US). BASS, Michael, B. [US/US]; 1743 North Marian
Avenue, Thousand Oaks, CA 91360-2163 (US).

(21) International Application Number: PCT/US01/08013

(74) Agent: NOONAN, Kevin; McDonnell Boehnen Hulbert
& Berghoff, 300 South Wacker Drive, Suite 3200, Chicago,
IL 60606 (US).

(22) International Filing Date: 13 March 2001 (13.03.2001)

(81) Designated States (*national*): AE, AL, AM, AT, AU, AZ,
BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE,
ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP,
KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD,
MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD,
SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ,
VN, YU, ZA, ZW.

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/188,786 13 March 2000 (13.03.2000) US

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,
CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(63) Related by continuation (CON) or continuation-in-part
(CIP) to earlier application:
US 60/188,786 (CON)
Filed on 13 March 2000 (13.03.2000)

Published:
— with international search report

(71) Applicant (*for all designated States except US*): AMGEN,
INC. [US/US]; One Amgen Center Drive, Thousand Oaks,
CA 91320-1799 (US).

(88) Date of publication of the international search report:
21 March 2002

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): JING, Shuqian
[CN/US]; 3254 Bordero Lane, Thousand Oaks, CA 91362

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.



WO 01/68854 A3

(54) Title: FIBROBLAST GROWTH FACTOR-LIKE MOLECULES AND USES THEREOF

(57) Abstract: The present invention provides Fibroblast Growth Factor-Like (FGF-L) polypeptides and nucleic acid molecules encoding the same. The invention also provides selective binding agents, vectors, host cells, and methods for producing FGF-L polypeptides. The invention further provides pharmaceutical compositions and methods for the diagnosis, treatment, amelioration, and/or prevention of diseases, disorders, and conditions associated with FGF-L polypeptides.

INTERNATIONAL SEARCH REPORT

Intern. Application No
PCT/US 01/08013

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C12N15/12 C07K14/50 C12N1/21 C12N5/10 C07K16/22
A61K39/395 G01N33/68 A61K38/18 A61K31/70 C12N15/62
A01K67/027 C12N5/22

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C12N C07K A61K G01N A01K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, EMBL, SEQUENCE SEARCH, CHEM ABS Data, BIOSIS, PAJ, WPI Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>KOGA CHIE ET AL: "Characterization of a novel member of the FGF family, XFGF-20, in <i>Xenopus laevis</i>" BIOCHEMICAL AND BIOPHYSICAL RESEARCH COMMUNICATIONS, ACADEMIC PRESS INC. ORLANDO, FL, US, vol. 261, no. 3, 11 August 1999 (1999-08-11), pages 756-765, XP002164263 ISSN: 0006-291X the whole document</p> <p style="text-align: center;">--- -/--</p>	1-11

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

A document defining the general state of the art which is not considered to be of particular relevance

E earlier document but published on or after the international filing date

L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

O document referring to an oral disclosure, use, exhibition or other means

P document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

G document member of the same patent family

Date of the actual completion of the international search

9 October 2001

Date of mailing of the international search report

24/10/2001

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Le Cornec, N

INTERNATIONAL SEARCH REPORT

Intern. Application No

PCT/US 01/08013

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>DATABASE EMBL 'Online! HS1150649, accession number AA232729, 6 March 1997 (1997-03-06) L. HILLIER ET AL: "WashU Merck EST project 1997. zr75g09.r1 Soares_NhHMPu Homo sapiens cDNA clone IMAGE:669280 5', mRNA sequence" XP002179649 * 99,6% identity in 496 nt overlap with seq ID no.1 (1-496:1-496) * abstract & UNPUBLISHED,</p>	1-3
X	<p>WO 96 41523 A (YAYON AVNER ;YEDA RES & DEV (IL)) 27 December 1996 (1996-12-27) claims; figure 2</p>	1-11,14, 17
A		12,13, 15-17, 37,47-54
X	<p>MIYAMOTO M. ET AL: "MOLECULAR CLONING OF A NOVEL CYTOKINE CDNA ENCODING THE NINTH MEMBER OF THE FIBROBLAST GROWTH FACTOR FAMILY, WHICH HAS A UNIQUE SECRETION PROPERTY" MOLECULAR AND CELLULAR BIOLOGY, WASHINGTON, DC, US, vol. 13, no. 7, 1 July 1993 (1993-07-01), pages 4251-4259, XP002071233 ISSN: 0270-7306</p>	1-12,14, 17,18, 20,23, 31,36
A	the whole document	15,19, 30,34, 39,44,50
X	<p>DATABASE EMBL 'Online! AB020858, accession number AB020858, 14 December 1998 (1998-12-14) M. HIRAKAWA ET AL: "Homo sapiens genomic DNA of 8p21.3-p22 anti-oncogene of hepatocellular colorectal and non-small cell lung cancer, segment 1/11" XP002164266 * 99,2% identity in 899nt overlap with seq ID no.1 (898-1:1592-16822) * abstract & UNPUBLISHED,</p>	1-3

-/--

INTERNATIONAL SEARCH REPORT

Intern. Application No

PCT/US 01/08013

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	SANTOS-OCAMPO ET AL: "Expression and biological activity of mouse fibroblast growth factor-9 (FGF-9)" JOURNAL OF BIOLOGICAL CHEMISTRY, AMERICAN SOCIETY OF BIOLOGICAL CHEMISTS, BALTIMORE, MD, US, vol. 271, no. 3, 19 January 1996 (1996-01-19), pages 1726-1731, XP002108162 ISSN: 0021-9258 the whole document	1-12,14, 17
A		15,19, 30,34, 39,44,50
P,X	HIROYUKI KIRIKOSHI ET AL: "Molecular cloning and characterization of human FGF-20 on chromosome 8p21.3-p22" BIOCHEMICAL AND BIOPHYSICAL RESEARCH COMMUNICATIONS, ACADEMIC PRESS INC. ORLANDO, FL, US, vol. 274, no. 2, 2 August 2000 (2000-08-02), pages 337-343, XP002164265 ISSN: 0006-291X * 100% identity in 854 nt overlap with seq ID no.1 (477-1330:1-854) * the whole document	1-17
P,X	WO 00 54813 A (DWARKI VARAVANI J ;RENDAHL KATHERINE (US); CHIRON CORP (US); MCGEE) 21 September 2000 (2000-09-21) claims figure 29	1-17,33, 37,42-44
P,X	WO 01 07595 A (BURGESS CATHERINE ;LICHENSTEIN HENRI (US); PRAYAGA SUDHIRDAS K (US) 1 February 2001 (2001-02-01) sequences ID no.1 and 2 claims; examples	1-54
E	WO 01 31008 A (UNIV KYOTO ;CHIRON CORP (US)) 3 May 2001 (2001-05-03) claims; figures 8,9 examples seq ID no.3	1-54

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.1

Although claims 33, 47 and 52 are directed to a method of treatment of the human/animal body (rule 39.1 IV PCT), the search has been carried out and based on the alleged effects of the compound/composition.

Although claim 36 (as far as it concerns an in vivo method) and claim 48 are directed to a diagnostic method practised on the human/animal body (rule 39.1 IV PCT), the search has been carried out and based on the alleged effects of the compound/composition.

Continuation of Box I.2

Claims Nos.: 18-19, 30-33, 36 all partially

Claims 18, 19, 30, 31 and 32 refer to a selective binding agent that specifically binds to the polypeptide FGF-L without giving a true technical characterization. In consequence, the scope of said claims is ambiguous and vague, and their subject-matter is not sufficiently disclosed and supported.

Claim 33 refers to a method for treating, preventing or ameliorating a FGF-L polypeptide-related disease by administering a selective binding agent without giving a true technical characterization. In consequence, the scope of said claims is ambiguous and vague, and their subject-matter is not sufficiently disclosed and supported.

Claim 36 refers to a method of detecting or quantitating the amount of FGF-L polypeptide using the selective binding agent without giving a true technical characterization. In consequence, the scope of said claims is ambiguous and vague, and their subject-matter is not sufficiently disclosed and supported.

No search can be carried out for such purely speculative claims whose wording is, in fact, a mere recitation of the results to be achieved. But a partial search has been carried out as far as the selective binding agent is an antibody to the polypeptide as mentioned in claims 20-29, 34-35.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 01/08013

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9641523	A	27-12-1996	AU 715996 B2	17-02-2000
			AU 6014496 A	09-01-1997
			AU 721773 B2	13-07-2000
			AU 6014596 A	09-01-1997
			CA 2223701 A1	27-12-1996
			CA 2224229 A1	27-12-1996
			CN 1195982 A	14-10-1998
			EP 0833620 A1	08-04-1998
			EP 0836380 A1	22-04-1998
			WO 9641620 A1	27-12-1996
			WO 9641523 A1	27-12-1996
			JP 11508358 T	21-07-1999
			JP 11507828 T	13-07-1999
WO 0054813	A	21-09-2000	AU 3755900 A	04-10-2000
			WO 0054813 A2	21-09-2000
WO 0107595	A	01-02-2001	AU 6495800 A	13-02-2001
			WO 0107595 A2	01-02-2001
WO 0131008	A	03-05-2001	AU 1101901 A	08-05-2001
			WO 0131008 A2	03-05-2001